

The *BeppoSAX* WFC X-ray source catalogue

F. Verrecchia¹, J. J. M. in 't Zand², P. Giommi¹, P. Santolamazza¹, S. Granata¹, J. J. Schuurmans² and L. A. Antonelli^{1,3}

¹ ASI Science Data Center (ASDC)*, c/o ESA-ESRIN, via Galileo Galilei, I-00044 Frascati, Rome, Italy

² Netherlands Institute for Space Research (SRON), Sorbonnelaan 2, 3584, CA Utrecht, The Netherlands

³ INAF – Osservatorio Astronomico di Roma (OAR), via Frascati 33, Monteporzio Catone, Rome, Italy

Received: 29 December 2006 / Accepted: 26 June 2007

ABSTRACT

Aims. We present the catalogue of X-ray sources detected by the two Wide Field Cameras (WFCs) in complete observations on board *BeppoSAX* during its 6 years of operational lifetime, between April 1996 and April 2002.

Methods. The *BeppoSAX* WFCs were coded mask instruments sensitive in the 2–28 keV energy band with a 40x40 square degree fields of view, pointing in opposite directions and perpendicularly to the *BeppoSAX* Narrow Field Instruments (NFI). The WFCs were usually operated simultaneously to NFI observations, each lasting up to several days. WFCs observed thus the entire sky several times with a typical sensitivity of 2 to 10 mCrab. A systematic analysis of all WFC observations in the *BeppoSAX* archive has been carried out using the latest post-mission release of the WFC analysis software and calibrations.

Results. The catalogue includes 253 distinct sources, obtained from a total sample of 8283 WFC detections. We describe the basic statistical properties of the sample and present a six-year history of two celestial calibration X-ray sources.

Key words. catalogs — X-rays: binaries — X-rays: galaxies — X-rays: general — X-rays: stars

1. Introduction

The *BeppoSAX* X-ray astronomy satellite (Boella et al. 1997), was a major program of the Italian Space Agency (ASI) with participation of the Netherlands Agency for Aerospace Programs (NIVR). The main scientific goal of the mission was to perform spectroscopic and timing studies of several classes of X-ray sources in a very broad energy band (0.1–300 keV). The satellite was launched on April 30, 1996 and observations were carried out until April 30, 2002.

The scientific payload comprised four Narrow Field Instruments (NFIs) and two Wide Field Cameras (WFCs) pointing in opposite directions from each other and perpendicular to the NFI (Boella et al. 1997; Jager et al. 1997). The WFC observations covered the whole sky multiple times over the period 1996–2002 (see Fig. 1).

The WFCs large field of view (FOV) and good sensitivity made these instruments suitable to detect and locate X-ray transient sources with a precision of a few arcminutes and to monitor large parts of the sky. Among the most important results obtained with the WFCs is the discovery of the X-ray afterglow phenomenon of Gamma Ray Bursts in 1997 (Costa et al. 1997). Moreover, the 12 monitoring campaigns of the Galactic Bulge allowed the spectral and timing monitoring of known persistent and transient X-ray sources and the discovery of many new galactic X-ray transients and phenomena (see for example Ubertini et al. 1999; Cornelisse et al. 2000; in 't Zand 2001 and 2004a; Cornelisse et al. 2003). These observations resulted in a significant advance in the knowledge of the transient emission properties of low mass X-ray binaries (e.g., in 't Zand et al. 2004b, 2004c).

After the end of the mission a systematic data analysis of the entire WFCs archive was carried out. This analysis, based on the processing of two complementary datasets, allowed us to build the complete catalogue of WFC X-ray sources and an archive of high-level data products, including energy spectra and light curves of the brighter sources. This archive is accessible on the web through the ASDC multi-mission interactive archive (<http://www.asdc.asi.it/mmia/>).

In this paper we present the final *BeppoSAX* WFC source catalogue including 253 distinct and previously known X-ray sources except for a new transient. The catalogue was obtained from a total sample of 8283 true detections.

This catalogue is the first complete all-sky survey above 2.5 keV since *HEAO-1 A4* (Levine et al. 1984) with 6 years of coverage corresponding to a net total exposure of over 100 million seconds and with a far better location accuracy. It contains more information on variability over long time scale in the hard X-ray band than for example the ROSAT all-sky survey catalogues, the ASCA 0.7–10 keV (Sugizaki et al. 2001) and the first three INTEGRAL-IBIS low galactic latitude surveys (Bird et al. 2004, Bird et al. 2006, Revnivtsev et al. 2004) and the high galactic latitude survey of the Swift-BAT (Markwardt et al. 2005), and comparable with the last INTEGRAL-IBIS survey (Bird et al. 2007).

2. Instrument and software description

2.1. General characteristics and the raw data archive

The WFCs were two identical coded mask instruments for the imaging of the hard X-ray sky (1.8–28 keV; Jager et al. 1997) developed and built by SRON. Their FOV was 40° × 40° each (full width to zero response) with an angular resolution of 5' (full width at half maximum, FWHM); the spectral and time resolution were 20% at 6 keV (FWHM) and 0.5 ms, respectively.

Send offprint requests to: verrecchia@asdc.asi.it

* INAF-OAR personnel resident at ASDC under ASI contract I/024/05




Fig. 1. The Aitoff projection in galactic coordinates of all the WFC observation pointing directions (left) and of the FOV of a typical WFC pointing of the Galactic Center (the circle includes the squared WFC FOV). WFC unit 1 points at Sgr-A and unit 2 at the anti-center, including the Crab Nebula.

The source location accuracy for bright sources (for the standard attitude reconstruction from star sensor data) was $1.4'$ at 99% confidence level, and larger for less significant sources as measured through the signal-to-noise ratio (SNR). The on-axis sensitivity was in the range 2–10 mCrab for a typical *BeppoSAX* WFC observation of 3×10^4 s, and depended on the intensities of other sources in the same FOV.

These properties made the WFCs suitable instruments for the monitoring of medium to high intensity X-ray sources and for the study of transient X-ray phenomena.

The WFCs raw data archive has been stored in 2652 Final Observation Tapes (FOTs). FOTs corresponding to contiguous observing periods with the same pointing and roll angle, were combined to form a multi-FOT complete observation.

We carried out two data processing runs on two different data samples. The first pertains to all data when no part of the FOV is occulted by the Earth. Some FOTs were, thus, completely discarded and 1438 observations remained. The second sample pertains to all data (1543 observations), whether occulted by the Earth or not, and exposure corrections were applied for each source to take into account occultations. We performed two different processing runs because the first allowed a straightforward modelling of the detector background image resulting in less image artifacts, while the second allowed to obtain the longest source exposure to reduce statistical noise and increase source time coverage.

2.2. The WFC imaging algorithm

The WFCs are shadow mask cameras (Ables 1968; Dicke 1968). An opaque screen with a pseudo-random pattern of holes (coded mask) is placed in front of a position-sensitive detector (a multi-wire proportional counter; Mels et al. 1988), so that X-ray sources in the FOV cast shadows on the detector, each displaced according to the off-axis position of the source. The WFC masks and detectors were of the same size and, therefore, only the on-axis position is fully coded with the mask pattern. However this configuration allows a better angular resolution at the same FOV compared to systems having the entire FOV fully coded.

The sky image is reconstructed with an iterative cleaning algorithm ('Iterative Removal Of Sources', IROS; Hammersley 1986, Hammersley et al. 1992; in 't Zand 1992), based on a cross-correlation of the detector image with the coded mask pat-

tern (Fenimore & Cannon 1978). IROS performs a new cross-correlation during each iteration and new detections are localised via a fit with the expected Point Spread Function (PSF) and then compared to the positions of X-ray sources in a reference catalogue. Finally, each source that is significantly above the noise, is simulated on the detector plane, including its energy and angle-dependent spatial response, and then subtracted.

The main systematic error in the reconstructed sky is due to a residual spatial non-linearity in the detectors which cannot be accounted for. As a result physical pixel sizes are not completely identical and consequently the PSF shows inaccuracies. The main impact is on the sensitivity in very crowded FOVs for exposures longer than a few days (e.g., the Galactic Center field, hereafter GC) and on the source location accuracy which is, in this case, hard limited to a best-case value of $1.4'$ (99% confidence level; Heise et al. 1998).

2.3. The reference catalogue

The reference catalogue of X-ray sources used in the first data processing run includes 17674 entries. It was built combining various tables of known astronomical objects (see Table 1) in order to include a large sample of entries from extra-galactic catalogues to allow the detection of AGN and cluster of galaxies in the 2–10 keV band at a flux level near the limiting sensitivity. Each WFC FOV includes from a few hundred up to two thousand sources of the reference catalogue. This requires an adjustment of the detection threshold criteria since source positions are more likely to coincide with a noise peak when there are many "potential" sources in the FOV. In the second data processing run a smaller reference catalogue was used to reduce the probability of a coincidence of a known source with a noise peak. It was built by selecting sources with 1 keV flux density greater than 4.8×10^{-12} erg cm $^{-2}$ s $^{-1}$ keV $^{-1}$ and including validated sources from the first processing run.

2.4. WFC processing procedure

Raw data consist of a list of events labelled with detector position, energy channel and detection time. Fluxes and positions of sources detected in an observation are determined from the raw data in three main steps (Jager et al. 1997):

- *I*: a 'cleaned' event list is obtained from each FOT for time intervals with known and stable attitude, excluding all un-

Table 1. The astronomical databases used to build the reference catalogue. Database acronyms, objects class and the references for each catalogue are shown from left to right. GC and GP are the Galactic Center and Plane.

Database Name	Objects class	References
XRB	X-ray Binary systems	van Paradijs 1995
HMXRB	High Mass X-ray Binary systems	Liu et al. 2000
LMXRB	Low Mass X-ray Binary systems	Liu et al. 2001
INTEGRAL-IBIS GP Soft γ -ray Catalogue	X-ray Binary systems	Bird et al. 2004
INTEGRAL-IBIS Hard X-ray GC Survey		Revnivtsev et al. 2004
Swift-BAT High Galactic Latitude survey	AGN	Markwardt et al. 2005
ZCAT	Galaxies with B magnitude < 14	Huchra et al. 1992
AGN	Active Galactic Nuclei	Padovani et al. 1997 and private communication
RASS-BSC	ROSAT All-Sky Survey Bright Sources with $0.1\text{--}2.4$ keV flux $> 10^{-11}$ erg s $^{-1}$ cm $^{-2}$	Voges et al. 1999
Sedentary	High Energy Peaked BL LACs (HBL)	Giommi et al. 1999, Giommi et al. 2005
CABSCAT	Spectroscopic binary systems	Strassmeier et al. 1988, Strassmeier et al. 1993
CVCAT	Cataclismic variable stars	Downes & Shara 1993
ALLWARPS	Cluster of galaxies	Perlman et al. 2002
BASSANI	Absorbed active galactic nuclei	private communication

wanted periods (such as those due to abnormal values of housekeeping parameters and passages through the South Atlantic Geomagnetic Anomaly). In the first processing run times were selected to keep the Earth outside the FOV, while in the second one periods were included when the Earth is inside the FOV (determining for each source the exposure when it is not occulted). Event lists from each FOT in a complete observation are merged;

- *II*: data products such as detector and sky images, and fluxes in specific energy bands are generated from event lists through the decoding process. A first decoding run is executed in the most sensitive band (i.e. 2–8 keV, assuming a Crab-like spectrum), to select significant sources. Then the decoding algorithm is executed in all energy bands for these sources. In the second processing run a further decoding test in the 8–19 keV band was added to search for hard transient sources;
- *III*: higher level scientific products, such as spectral fits with simple models, spectral energy distributions and light curves are generated in the same bands for each detected source, applying SNR acceptance thresholds of 4.5 and 6.5 for sparse and crowded fields respectively.

The SNR is calculated by the WFC software using as noise the standard deviation evaluated on source plus background counts. This is because for the most part of the sources the standard deviation is background dominated. Thresholds were preliminarily defined taking into account the large reference catalogue and the sensitivity dependence on the combined intensity of all sources in the FOV (see detailed description in section 3.3).

At the end of the standard procedure, a visual inspection of the scientific products (spectra and light curves) was performed.

3. Results

In this section we describe how the fluxes were validated with data of well-documented stable X-ray sources and how the WFC catalogue was built through the creation of the final sample of detections.

3.1. The observations of the Crab Nebula and Cas A

As validation of both the WFC software and the standard analysis procedure at high, medium and low flux levels, we checked

all the observations of a sample of well-documented stable X-ray sources including the Crab Nebula, Cas A and the Tycho supernova remnant. We show here some results for the Crab and Cas A from the first data processing run.

We analyzed 130 observations of the Crab Nebula at off-axis angles lower than 21° and exposures ranging between 1.67 and 129.7 ks. We calculated the relative deviations from the Crab Nebula 2–10 keV mean flux (2.12×10^{-8} erg cm $^{-2}$ s $^{-1}$, evaluated using the final WFC response matrices) and corrected fluxes for a residual linear trend with off-axis angle between FOV center and borders of about 7%. This trend is mainly due to the decrease with the cosine of the angle of the number of photons detected per square cm. We plotted then the resulting 2–10 keV secular light curve of Crab Nebula (Fig. 2). A systematic error of about 2% was added in quadrature to the flux errors to make the data consistent with an expected reduced chi-square of 1. This error was applied to all the detections in the catalogue. We show in the bottom panel of Fig. 2 the Cas A corrected secular light curve obtained from 144 observations, compatible with a residual dispersion of $\sim 6.2\%$. This dispersion is larger than the Crab one at the lower flux level of Cas A. A further systematic error of about 4.1% is needed to obtain again a reduced chi-square of 1. This term has not been added to flux errors.

Table 2. Number of sources and detections for the main source classes in the WFC catalogue. The table does not include 37 detections of 2 X-ray pulsar sources, 4U 0142+614 and PSR J0540-6919.

Source Class	Number of detections	Number of sources
candidate X-ray binaries	2	2
High Mass X-ray binaries	2045	49
Low Mass X-ray binaries	4876	87
Cataclismic Variables	125	11
Extended Galactic sources	443	5
Stars	99	21
White Dwarfs	15	2
Galaxies	14	2
AGN	472	45
Cluster of Galaxies	59	13
Unclassified	96	14




Fig. 2. The 2-10 keV Crab Nebula (top panel) and Cas A secular light curves corrected for the linear decrease with off-axis angle described in the text (open circles correspond to unit 1). A systematic error of $\sim 2\%$ was added in quadrature to the corrected flux errors for both WFCs to reach a reduced chi-square of ~ 1 . Residual dispersions for corrected light curves are 2% and 6.2% respectively.

3.2. Source identifications

Sources in the reference catalogue (see section 2.3) were automatically identified by the IROS algorithm and then double-checked visually. Sources not identified or misidentified in the reference catalogue were cross checked with literature regarding sources discovered by ASCA, RXTE, *BeppoSAX*, INTEGRAL, Swift-BAT, XMM (for example see Sugizaki et al. 2001, Giommi et al. 2000, Bird et al. 2006 and 2007, Markwardt et al. 2005, XMM-SSC 2003 and 2006). Moreover we checked and updated, when necessary, source classifications.

3.3. Source detection filtering

The sensitivity at a certain position in the FOV depends on the off-axis angle (due to a dependence on effective area), on the particle-induced and cosmic diffuse backgrounds and on the combined intensity of all X-ray point sources in the FOV. The first three components are rather stable under nominal conditions, the latter varies considerably with galactic coordinates

(the variations of the cosmic background above 2 keV are much smaller; see for example Kushino et al. 2002 and Lump et al. 2002). The largest concentration of bright X-ray sources is close to the GC, where consequently the sensitivity is worse than elsewhere in the sky.

In fields crowded with bright sources there is a complicating factor related to a non-optimum imaging by the detectors. The detectors suffer from a differential non-linearity (Jager et al. 1997). Due to the dependence on the spectrum and off-axis angles of all detected sources, it is not completely accounted for in the IROS algorithm. The effect is so strong that a systematic sensitivity limit is reached for GC fields for an exposure longer than a few hundred ks.

In order to distinguish those fields where the contribution to the background from bright sources is dominant, we considered the Total Count Rate (hereafter TCR) of an observation. TCR is defined as the total number of events divided by the net exposure time. Using this parameter we selected observations of low background fields ($\text{TCR} < 90 \text{ cts s}^{-1}$ and $\text{TCR} < 130 \text{ cts s}^{-1}$ in the first and second data processing run respectively¹), such as fields at high galactic latitude or those not crowded, and the high background ones at low latitude ($\text{TCR} > 90, 130 \text{ cts s}^{-1}$).

The criterion to accept the detection of a point source is usually different between known and new sources. The acceptance threshold of a known source at a certain position is defined as the level of 3σ excess above the statistical noise induced at that position by all contributions mentioned above (where σ is the statistical standard deviation). This corresponds, for a gaussian noise distribution, to a chance probability of less than 0.14%. Since in a coded aperture camera the flux for every position in the sky is reconstructed from photons on large portions of the detector, the noise distribution is still Gaussian.

In the case of a previously uncatalogued X-ray source, its position is not known so one needs to take into account the number of trials. This number is firstly determined from the number of independent pixels in the FOV, 255×255 (i.e., the WFC number of mask elements). If we require less than 1 false peak in 100 images, the threshold needs to be 5.3σ .

In our first procedure we implemented a very large reference catalogue, which produces in almost all WFC FOVs a large number of “potentially” visible known sources (see section 2.3). This increases proportionally the probability for false detections (1000 potential sources in the FOV imply about 2 sources coincident by chance with 3σ excesses). This is not completely true however for GC fields, where the detectors non-linearity requires a fine tuning of the detection acceptance threshold.

In order to take into account all the factors and evaluate the required significance threshold, we built the cumulative distribution of all the 2–8 keV detection SNRs (Fig. 3) in both data processing runs. This plot is roughly similar to a “logN-logS” distribution.

We fit the data to a broken power law plus a cumulative gaussian model (BPL + CG) and in Fig. 3 (left panel) we show both the components resulting from a fit to data from the first processing run stage (CG has a mean of 3.9 and $\sigma = 1.2$, while the BPL has slopes of ~ -0.53 , ~ -0.01 and a break at $\text{SNR} \sim 4.0$). A BPL component was used to distinguish better the statistical

¹ In the second processing run sources have different net exposures, so we built this parameter using the shortest common exposure time among them. We verified these ranges of values using the “root mean squares” of significance images discussed below in this section, as an alternative test of a field background.




Fig. 3. The cumulative 2–8 keV SNR distributions of all the detections without any SNR cut for the first data processing run (left panel) and for the most complete one. The upper black thick curves are the total samples, the dash-dotted curves are the components of the fitted models (Gaussian + Broken power law, with parameters mean = 3.9, $\sigma = 1.2$, slopes ~ -0.53 and ~ -0.01 and break at ~ 4.0 for the first processing run and mean = 4.0, $\sigma = 1.2$, same slopes and break at ~ 5.8), while the remaining three lower curves are the distributions for different off-axis angle intervals (the circled one corresponds to off-axis angles $< 10^\circ$, the squared to $10\text{--}15^\circ$ and the triangled to $> 15^\circ$).

noise component, represented by the CG component, from the true detections, represented by the BPL component. The BPL takes into account the instrumental sensitivity limits, the break roughly indicating the significance above which we can expect “true” detections following a power law distribution.

The CG component, describing the background noise distribution (both random and systematic), has parameters which imply a sample of $\sim 4\%$ of spurious detections at $\text{SNR} \geq 6.3$. We visually checked each marginal detection (with $\text{SNR} < 9$, a distance from the reference position greater than $5'$ and at FOV borders). The percentage of discarded detections is $\sim 2.8\%$, leaving about 1.2% of residual noise detections in the first processing run sample.

In Fig. 3 we also show the cumulative distributions for detections in three off-axis angle intervals, showing that the highest contribution to the CG noise component comes from detections at off-axis angles larger than 15° ($\sim 53\%$ of all detections in the gaussian component). These plots confirm the results of the visual inspection. Selecting detections with SNR greater than 6 from the first processing run, we obtained 7295 detections of 228 distinct sources.

In the right panel of Fig. 3 we show the 2–8 keV SNR cumulative curves for the second processing run. These curves show a much lower CG component in all curves, evident only for off-axis angles larger than 15° . It was very difficult to estimate this component from the total curve. However the estimated CG parameters (mean ~ 3.6 and $\sigma \sim 1.2$) ensure a contamination from spurious detections ($\sim 1\%$ at SNR above 5.9) slightly larger than that obtained fitting the same component on the curve for off-axis angles larger than 15° . This lower CG component however confirms the effect of the reference catalogue on the first processing run.

In both processing runs the fitting process to the cumulative curves is not perfect. This is mainly due to the superposition of different noise components coming from different pointings (i.e., mainly from high/low background fields), but also to differences of all the other parameters which may influence

the sensitivity among observations (such as different exposure times). Therefore, we first checked the SNR distributions for the low TCR sub-samples for both processing stages. We estimated a gaussian component (mean ~ 3.0 and $\sigma \sim 1.0$) for the second stage which permits to establish a 1% contamination threshold in this curve at $\text{SNR} = 4.8$, while this component for the first processing run curve is comparable to that fitted on the total sample (threshold at 5.9).

As a final verification of the acceptability of low SNR (below $\text{SNR} = 7$) detections in low TCR fields, we computed specific 3σ confidence level thresholds for each field estimating their background noise standard deviation², and checked visually detections above these thresholds. Most of the low SNR detections in low background fields were discarded with this check, up to “single field” confidence levels of $\sim 7\sigma$.

We checked also results in harder bands from both processing runs, creating the same cumulative significance distributions (in Fig. 4 those relative to the second processing run). In particular for the 8–19 keV band we evaluated the CG component parameters (mean = 2.0 and $\sigma = 2.7$) using the same fit model, estimating a contribution of $\sim 2.5\%$ of noise detections above $\text{SNR} = 7$.

The catalogue has been obtained choosing detections from the second processing run having $\text{SNR} > 6$ if detected in the high TCR fields, or $\text{SNR} > 5$ if in the low ones. Finally we added to the catalogue a small sample of verified sources from the first processing run and not detected in the second one together with a sample of detections from harder bands. The complete sample of detections includes 8283 entries of 253 distinct sources.

² We evaluated these thresholds on significance images (the ratio of the sky image to the poisson noise map from the first iteration of IROS), either calculating the “root mean square” of a complete image and of sub-images (85×85 pixels) around each source, or estimating directly the gaussian background noise standard deviations. We corrected the 3σ levels for the number of potential sources in each FOV.




Fig. 4. The cumulative distributions of SNR in three larger/harder bands (2–26, 8–19, 19–26 keV) for detections of the second processing run. The gaussian component of the model fitted to the 8–19 keV curve is shown.

4. The Catalogue

The number of sources for the main classes of the 253 sources included in the catalogue are shown in Table 2. The positions and fluxes of the most significant detection of each source are reported in Table 3 (also available on-line at <http://www.asdc.asi.it/wfccat/>). At the end of the table a group of 11 rows includes sources detected in the first processing run only.




Fig. 5. The Aitoff projection of the *BeppoSAX* WFC detections: 8283 detections of 253 distinct X-ray sources in all the observations processed (symbol size is proportional to intensity)

Table 3 gives as first column the *BeppoSAX* WFC designation (i.e., “SAXWFC”), followed by right ascension and declination (all celestial coordinates are for epoch J2000). In the following column the positional error is presented, which was calculated by a quadratic sum of the 1.3’ systematic uncertainty (see below) and the mean of the 99% confidence level errors in X and Y. Then are presented the source net exposure (in ks), the 2–10 keV flux and error (in $\text{erg cm}^{-2} \text{s}^{-1}$) for the maximum SNR detection, followed by the detection SNR in 2–8 and 8–19 keV bands. The number of detections and the total exposure of each source are in the ninth and tenth column. The mean flux and its error (in $\text{erg cm}^{-2} \text{s}^{-1}$), the reduced chi-square and the minimum




Fig. 6. The exposure distributions of the *BeppoSAX* WFC detections in all data processed.

and maximum flux among all the detections of a source are in columns from the eleventh to the fourteenth. The last two columns contain the source identification and the classification resulting from comparison with public databases.

Some statistical properties of the catalogue are hereafter described. The Aitoff projection in galactic coordinates of the full sample is shown in Fig. 5. The observation exposure distributions (Fig. 6) have a peak at ~ 30 ks and ~ 50 ks for the first and second processing stages respectively, while only a few long high galactic latitude and GC pointings have exposures up to 350 ks in the second processing run. The total exposure is ~ 67 Ms for the first processing run while the second one has a minimum total exposure of ~ 102 Ms (choosing the minimum source exposure among those in a field, for each field). The catalogue mostly includes sources detected in the second processing stage.

The 2–10 keV flux vs the exposures for all the detections in the sample in Fig. 7 (top panel) shows up the flux limit reached ($\sim 3 \times 10^{-11} \text{ erg cm}^{-2} \text{s}^{-1}$) and the limiting sensitivity of the catalogue as a function of the exposure which follows the usual relation $F_x^{\text{limit}}(2-10 \text{ keV}) \approx c/T^{1/2}$, where the constant c is

$$c \approx \begin{cases} 7 \times 10^{-9} \text{ erg cm}^{-2} \text{s}^{-1/2} & \text{for } |b| > 30^\circ \\ 3 \times 10^{-8} \text{ erg cm}^{-2} \text{s}^{-1/2} & \text{for GC region} \end{cases} \quad (1)$$

These numbers are consistent with the predictions in Jager et al. 1997. It should be noted that for GC fields the sensitivity reaches a limit at about 100 ks of net exposure due to the above mentioned imperfect imaging. In the bottom panel of the same figure the 2–10 keV flux as a function of the off-axis-angle is given. An empirical model for the WFC limiting sensitivity, $F_x^{\text{limit}}(2-10 \text{ keV}) \approx (9.5 \times 10^{-16} \times R^{3.9} + 2.3 \times 10^{-11}) \text{ erg cm}^{-2} \text{s}^{-1} \text{deg}^{-1}$, where R is the off-axis angle, has been evaluated for high galactic latitude fields.

The location accuracy reached is shown in Fig. 8. Source location accuracy is known to depend on source SNR (in ‘t Zand 1992), so we plot the binned detection mean 99% position errors as a function of the SNR (top panel). In each bin we calculate the error average value and its standard deviation over detections having similar SNRs. No systematic error appears at




Fig. 7. The 2–10 keV flux as a function of the exposure (top panel) of the final sample of detections. The dashed lines on the bottom part of the plot show the limiting sensitivity model (proportional to the exposure square root) for fields out of the Galactic Plane (the lowest) and the GC ones. In the bottom panel the same flux vs the off-axis angle. The dashed curve on the bottom part of the plot shows a model of the limiting sensitivity as a function of the off-axis angle.

high SNRs. We show in the bottom panel the cumulative distributions of distances between measured and catalogued positions, for all the detections and for those with $\text{SNR} > 60$ and $\text{SNR} > 180$. Distances were obtained cross-correlating our list with the reference catalogue using a radius of $5'$. We computed then radii including 99% of the detections for all the distributions obtaining $3.30'$, $1.87'$ and $1.78'$ for the total sample and the two sub-samples at increasing SNR thresholds respectively. We fitted these curves with a power law plus cumulative gaussian model to evaluate the 99% confidence level error radius on the gaussian component, which is the expected distribution for bright “good” detections. The three curves illustrate how at higher SNR thresholds the distributions are better represented by the gaussian component. The radii for the three gaussian components are $1.78'$, $1.61'$ and $1.33'$ respectively. This last value is consistent with what previously reported for bright sources (Piro et al. 1997, Heise et al. 1998). This minimum location accuracy value has been added in quadrature as a systematic error to the mean 99% position error for each detection. The total po-




Fig. 8. Positional 99% error radius (in arcmin) as a function of the SNR (top panel), where each data point represents the average over 150 source positions and the standard deviation with respect to that average. In the bottom panel the cumulative distribution of the distances between measured positions and those in the reference catalogue (in arcminutes) for all the detections (upper curve) and for detections with $\text{SNR} > 60$ and $\text{SNR} > 180$. The radii including 99% of the detections are $\sim 3.3'$, $1.87'$ and $1.78'$ respectively. The dashed curve is the cumulative gaussian component of the fit model applied to the upper curve, which represent the expected model for bright sources.

sition error computed for the maximum SNR detection of each source is reported in Table 3.

As an example of the location accuracy for a bright source we show in Fig. 9 the positions of all Cyg X-1 detections in equatorial coordinates. We fitted the distribution of Cyg X-1 positions in both coordinates with a gaussian model obtaining a best fit mean position, R.A.= 299.5898, Dec.= 35.2021 (J2000) with a $\sim 1.6'$ 95% confidence level radius.

The chance coincidence probability of the correlation between WFC detections and the reference catalogue positions used in the source post-processing identification, can be evaluated by shifting the detection positions by various fixed amounts (between -10° and 10°) and executing again the cross-correlation with the same radius (see Fig. 10). The number of spurious matches for shift values greater than 0.16° is below $\sim 3\%$.

We calculated two hardness ratios (HR1, HR2) using the 1.7–3.2, 4.6–7.1 and 7.1–10.9 keV bands for two sub-samples of 7439 and 7529 detections respectively, having non-zero values of count rates in these bands. The HR1 distribution for each individual source class included in the catalogue (see Table 2) is shown in Fig. 11. These histograms include HR1 values from all possible sources states, such as hard/soft states for LMXRBs and HMXRBs, all overlapped. In particular for Supergiant plus unclassified HMXRBs HR1 covers a very broad range of values with two peaks at ~ 0.6 and ~ 3 .

The HR2 distributions for each source class are shown in the left panel of Fig. 11. In this case the plot for Supergiant HMXRBs class shows a prevailing harder peak at ~ 1 , where the Be HMXRBs distribution peaks too. Two energy indexes were evaluated for both HRs assuming galactic absorption values for all sources. They are shown in the two panels of the Fig. as top scales. Their values range between -0.6 and 3.5 for that evaluated from HR1 and between 0.5 and 3.0 for the harder one.

5. Conclusions

We presented the complete post-mission catalogue of WFC X-ray sources detectable in complete observations, which includes 253 distinct objects. As shown in Table 2, these sources are mainly galactic (176 sources plus three candidate galactic transient sources, i.e. 71% of the total), while the smaller extragalactic sample, 62 sources (25%), is mainly composed of AGN, 11 of which are BL Lacs (4.3%). The galactic sources are mostly LMXRBs (48%) while the HMXRBs (27%) include 13 classified Supergiant systems (7.3%) and 23 Be HMXRBs (13%). The 14 unclassified sources in Table 3, include two known sources, SAX J1428.6-5422 and SAX J1805.5-2031, and a new transient X-ray source, for which the WFC detections give the best lo-




Fig. 9. The positions of all Cyg X-1 detections in equatorial coordinates. Crosses correspond to WFC unit 1, circles to unit 2, while the star corresponds to the Cyg X-1 reference catalogue position, R.A.= 299.5904, Dec.= 35.2016 and the encircled cross to the best fit position, R.A.= 299.5898, Dec.= 35.2021. The thick circle of radius $\sim 1.6'$ represents the 95% confidence level centered on the gaussian model best fit position.




Fig. 10. Spurious matches in the cross-correlation of the *BeppoSAX* WFC catalogue with the reference catalogue with a radius of $5'$ for “shift” values from 0° to 2° .

calization so far. These are candidate transient galactic sources detected in outburst. Also included is a more uncertain candidate X-ray binary source (RX J0121.4-7258) and a possible BL Lac object (1RXS J205528.2-002123).

Most of the catalogue sources (83%) have a counterpart in the softer RASS catalogue (within $3'$ radius). Moreover large fractions of sources have significant ($\text{SNR} \geq 7$) detections in the 2-5.5, 2-19 and 8-19 keV bands (219, 197 and 133 sources respectively), which have slightly lower percentages (71%, 70% and 64% respectively) of counterparts in the RASS. On the other hand the correlation with the harder INTEGRAL/IBIS Soft γ -ray catalogues (Bird et al. 2004, 2006 and 2007) give increasing percentages proportionally to the increasing INTEGRAL total exposure and sky coverage (29% with the first, 40% with the second and 53% with the most recent one, using the same $3'$ radius). If we consider a sample of harder WFC sources (35%), detected ($\text{SNR} \geq 10$) in the 11-26 keV band, 78% resulted to have a counterpart in the last IBIS catalogue. The WFC catalogue thus includes sources detected up to 19 keV with still a good sensitivity (sources detected in this band have a 2–10 keV flux greater than about 3 mCrabs).

The cross-correlation with the RXTE “All Sky Monitor” source catalogue (available on-line³) finally gives the highest number of results (75%) among catalogues in hard X-ray bands. This result confirms the importance of the WFC source catalogue as a complementary reference catalogue for current and future hard X-ray missions, particularly in the 2–10 keV band.

Acknowledgements. The ASDC is supported by the Italian Space Agency (ASI). Part of this work benefitted of ASI grant I/024/05. FV, PG and PS thank S. Piranomonte for the help to the revision of source identifications and classes detected in the first data processing run. Authors thank all people of the WFC team at SRON, particularly G. Wiersma, for his collaboration and kindness.

References

- Abell, G.O., Corwin, H.G.Jr., & Olowin, R.P. 1989, *ApJS*, 70, 1
- Ables, J.G. 1968, *Proceedings of the Astronomical Society of Australia*, 1, 172
- Perlman, E.S., Horner, D.J., Jones, L.R., et al. 2002, *ApJS*, 140, 265
- Bird, A.J., Barlow, E.J., Bassani, L., et al. 2004, *ApJ*, 607L, 33
- Bird, A.J., Barlow, E.J., Bassani, L., et al. 2006, *ApJ*, 636, 765

³ <http://xte.mit.edu/asmlc/asmcat.html>




Fig. 11. The Hardness Ratio 1 (1.7–3.2/4.6–7.1 keV bands; left panel) distributions for each source class included in the catalogue. On the right the same per classes distributions of Hardness Ratio 2 (4.6–7.1/7.1–10.9 keV). Two corresponding energy indexes, evaluated assuming galactic absorption, are shown in the top scales of both panels. Numbers below each class label, in both panels, are the detections having non-null values of count rates and of energy indexes.

- Bird, A.J., Malizia, A., Bazzano, A., et al. 2007, *ApJS*, 170, 175
Boella, G., Butler, R.C., Perola, G.C., et al. 1997, *A&AS*, 122, 299
Cornelis, R., Heise, J., Kuulkers, E., Verbunt, F., in 't Zand, J.J.M. 2000, *A&A*, 357, L21
Cornelis, R., in 't Zand, J.J.M., Verbunt, F., et al. 2003, *A&A*, 405, 1033
Costa, E., Frontera, F., Heise, J., et al. 1997, *Nature*, 387, 783
Dicke, R.H. 1968, *ApJ*, 153, L101
Downes, R.A. & Shara, M.M. 1993, *PASP*, 105, 127
Fenimore, E.E., Cannon, T.M. 1978, *ApJ*, 17, 337
Giommi, P., Menna, M.T., Padovani, P. 1999, *MNRAS*, 310, 465
Giommi, P., Perri, M., Fiore, F. 2000, *A&A*, 362, 799
Giommi, P., Piranomonte, S., Perri, M., Padovani, P. 2005, *A&A*, 434, 385
Hammersley, A. 1986, Ph.D. thesis, University of Birmingham
Hammersley, A., Ponman, T., Skinner, G.K. 1992, *Nucl. Instr. Meth. Phys. Res.* A311, 585
Heise, J., in 't Zand, J.J.M., Muller, et al. 1998, in "4th Huntsville Symp. on Gamma-Ray Bursts", ed. C. Meegan, R. Preece, & T. Koshut (New York: AIP), AIP Conf. Proc. 428, 397
Huchra, J.P., Geller, M.J., Clemens, C.M., Tokarz, S.P., Michel, A. 1992, *Bull. Inf. Centre Donnees Stellaires* 41, 31
Jager, R., Mels, W., Brinkman, A., et al. 1997, *A&AS*, 125, 557
Kushino, A., Aoki, Y., Yamasaki, N.Y., et al. 2002, *PASJ*, 54, 327
Levine, A.M., Lang, F.L., Lewin, W.H.G., et al. 1984, *ApJS*, 54, 581
Liu Q.Z., van Paradijs, J., van den Heuvel, E.P.J. 2000, *A&AS*, 147, 25
Liu Q.Z., van Paradijs, J., van den Heuvel, E.P.J. 2001, *A&AS*, 368, 1021
Lumb, D.H., Warwick, R.S., Page, M., De Luca, A. 2002, *A&A*, 389, 93
Markwardt, C. B., Barbier, L., Barthelmy, S., et al. 2005, *A&AS*, 207, 3801
Mels, W. A., Lowes, P., Buurmans, H. B., et al. 1988, *Nucl. Instr. Meth. Phys. Res.*, A273, 689
Padovani, P., Giommi, P., Fiore, F. 1997, *Mem. Soc. Astron. Ital.* 68, 147
Piro, L., Butler, R.C., Fiore, F., et al. 1997, "Addendum to the SAX Observers' Handbook", on-line Techn. Report, <http://www.asdc.asi.it/bepposax/handbook/>
Revnivtsev, M.G., Sunyaev, R.A., Varshalovich, D.A., et al. 2004, *PAZh*, 30, 430
Strassmeier, K.G., Hall, D.S., Zeilik, M., et al. 1988, *A&AS*, 72, 291
Strassmeier, K.G., Hall, D.S., Fekel, F.C., & Scheck, M. 1993, *A&AS*, 100, 173
Sugizaki, M., Mitsuda, K., Kaneda, H., et al. 2001, *ApJS*, 134, 77
Ubertini, P., Bazzano, A., Cocchi, M., et al. 1999, *ApJ*, 514, L27
van Paradijs, J. 1995, in "X-Ray Binaries", eds. Lewin W.H.G., Van Paradijs J., Van Den Heuvel E. P.J., Cambridge University Press.
Voges, W., Aschenbach, B., Boller, Th., et al. 1999, *A&A*, 349, 389
in 't Zand, J.J.M. 1992, Ph.D. thesis, University Utrecht
in 't Zand, J.J.M. 2001, in 'Exploring the gamma-ray universe', ed. A. Gimenez, V. Reglero, & C. Winkler, ESA Pub. Division, 163
in 't Zand, J.J.M., Verbunt, F., Heise, J., et al., 2004a, *NuPhS*, 132, 486
in 't Zand, J.J.M., Cornelisse, R., & Cumming, A. 2004b, *A&A*, 426, 257
in 't Zand, J.J.M., Cornelisse, R., Kuulkers, E., Verbunt, F., & Heise, J. 2004c, in "X-Ray Timing 2003: Rossi and Beyond", ed. P. Kaaret, F.K. Lamb, & J.H. Swank (Melville, NY: AIP), AIP Conf. Proc., 714, 253
Xmm-Newton Survey Science Centre, Consortium, 2003, *yCat.*, 9037, 0
The Second XMM-Newton Serendipitous Source Pre-release Catalogue, XMM-Newton Survey Science Centre, 2006.

BeppoSAX WFC Catalogue

Supplementary Table (Table 3)

Table 3. BeppoSAX WFC X-ray Source Catalogue

WFC Name	RA(2000) hh mm ss	DEC(2000) ° ' "	$\sigma_{ERR_{RAD}}$ arcmin	EXPO ks	^{b} Flux $_{2-10}$	^{c} SNR1	^{d} SNR2	N $_{det}$	EXPO $_{total}$ ks	^{e} <FLUX> $_{2-10}$	χ^2_{flux}	^{f} Flux $_{MAX}$	^{f} Flux $_{MIN}$	Name	^{g} Class
SAXWFC J0025.5+6407.3	00 25 28.7	64 07 18.8	1.3	74.0	17.6 ± 0.8	24.9	1.5	105	6358.6	17.1 ± 2.9	3.5	28.0	11.0	Tycho	SNR shell
SAXWFC J0028.7+5917.2	00 28 44.7	59 17 11.7	3.5	45.3	4.7 ± 0.6	7.8	2.8	9	760.2	5.1 ± 1.3	1.6	8.6	3.9	V709 Cas	CV
SAXWFC J0035.9+5949.7	00 35 52.2	59 49 42.2	1.5	101.2	8.2 ± 0.7	12.7	3.1	10	809.3	6.6 ± 1.9	4.5	11.0	4.5	1ES 0033+595	AGN, BL Lac
SAXWFC J0037.0+6121.0	00 37 01.7	61 21 02.8	1.4	46.4	12.0 ± 0.7	16.9	5.4	6	357.6	9.6 ± 7.6	27.9	27.0	6.0	IGR J00370+6122	HMXRB
SAXWFC J0051.8-7217.5	00 51 45.0	-72 17 31.2	1.5	33.9	9.3 ± 0.6	15.8	6.4	1	—	—	—	—	—	SMC X-3	HMXRB, XP
SAXWFC J0052.3-7319.3	00 52 20.1	-73 19 17.0	1.3	169.5	15.6 ± 0.6	26.8	10.7	12	571.1	17.0 ± 7.3	23.8	35.0	6.2	RX J0052-7319	Be HMXRB, XP,
SAXWFC J0053.2-7438.8	00 53 09.5	-74 38 50.2	1.6	109.8	6.3 ± 0.6	11.1	5.4	2	174.7	6.6 ± 0.5	0.8	7.1	6.3	CF Tuc	RSCVn
SAXWFC J0054.3-7340.2	00 54 19.7	-73 40 12.7	2.1	76.6	13.1 ± 2.0	5.7	4.0	2	240.9	10.2 ± 3.0	3.0	13.0	8.8	SMC X-2	Be HMXRB, XP,
SAXWFC J0056.7+6043.4	00 56 44.9	60 43 26.7	1.3	248.0	16.3 ± 0.6	32.3	11.1	72	4321.3	14.2 ± 3.6	9.2	24.0	7.7	Gamma Cas	Be HMXRB
SAXWFC J0111.3-7317.2	01 11 18.1	-73 17 13.2	1.3	140.0	57.2 ± 1.2	124.2	59.3	17	1083.6	17.4 ± 20.5	300.6	62.0	4.8	XTE J0111-7317	Be HMXRB, XP,
SAXWFC J0117.0-7326.4	01 17 00.5	-73 26 22.5	1.3	103.8	81.5 ± 1.7	114.3	61.0	158	11287.8	39.6 ± 26.2	282.6	127.0	5.5	SMC X-1	SG HMXRB, XP
SAXWFC J0118.0+6517.7	01 18 02.3	65 17 42.7	1.3	39.5	28.3 ± 0.7	57.7	44.9	48	2978.6	14.2 ± 7.6	46.9	39.0	7.3	2S 0114+65	SG HMXRB, XP
SAXWFC J0118.5+6343.8	01 18 29.8	63 43 47.2	1.3	108.3	163.5 ± 3.6	79.2	56.1	10	535.3	31.9 ± 126.2	450.1	420.0	10.0	4U 0115+634	Be HMXRB, XP,
SAXWFC J0121.7-7257.6	01 21 41.0	-72 57 33.4	1.4	64.9	16.6 ± 0.7	24.1	19.7	2	157.7	17.5 ± 2.7	6.6	20.0	17.0	RX J0121.4-7258	unclassified
SAXWFC J0124.0-3504.6	01 23 57.0	-35 04 36.1	2.0	69.1	6.2 ± 0.6	9.5	4.0	3	141.8	7.1 ± 6.5	12.3	18.0	6.2	NGC 526A	AGN, Seyfert 1
SAXWFC J0146.5+6145.3	01 46 32.2	61 45 15.4	1.5	259.5	8.3 ± 0.6	13.5	0.8	31	2368.4	7.7 ± 2.0	3.2	16.0	5.5	4U 0142+614	AXP
SAXWFC J0147.0+6120.6	01 47 01.2	61 20 34.0	1.5	81.9	9.8 ± 0.7	14.0	6.5	23	1875.3	7.7 ± 5.2	8.9	31.0	4.2	RX J0146.9+6121	Be HMXRB, XP
SAXWFC J0209.5+5226.2	02 09 31.8	52 26 13.2	2.3	59.0	15.9 ± 2.7	5.1	0.3	1	—	—	—	—	—	1H 0203+513	AGN, Seyfert 1
SAXWFC J0245.0+6225.7	02 44 57.5	62 25 42.6	1.7	38.0	6.5 ± 0.7	9.7	3.6	7	535.6	5.1 ± 0.7	1.2	6.5	4.2	QSO B0241+62	AGN, Seyfert 1
SAXWFC J0254.3+4133.3	02 54 20.7	41 33 17.2	1.9	103.4	5.9 ± 0.9	6.6	1.5	3	333.1	5.2 ± 0.6	0.5	5.9	4.7	2A 0251+413	CIG
SAXWFC J0308.2+4057.1	03 08 11.9	40 57 08.6	1.3	20.9	68.9 ± 2.4	31.0	2.8	6	241.3	11.5 ± 23.7	165.3	69.0	4.1	Algol	Algol star type
SAXWFC J0311.3-2046.2	03 11 20.5	-20 46 09.1	2.1	14.4	12.7 ± 2.0	6.0	0.0	1	—	—	—	—	—	HE 0309-2057	AGN, Seyfert 1
SAXWFC J0318.0-4414.7	03 18 00.0	-44 14 44.5	2.1	82.7	5.6 ± 0.8	7.0	4.2	3	328.9	3.5 ± 1.6	5.2	5.6	2.8	ABELL 3112	CIG
SAXWFC J0319.8+4131.0	03 19 47.1	41 31 01.2	1.3	25.0	40.6 ± 1.6	29.5	6.0	40	1377.8	37.3 ± 3.7	2.6	49.0	32.0	NGC 1275	AGN, RG
SAXWFC J0326.6+2842.7	03 26 33.0	28 42 44.2	1.5	81.9	9.9 ± 0.6	15.5	3.8	2	157.6	9.3 ± 1.5	3.1	9.9	7.9	UX Ari	RSCVn
SAXWFC J0331.3+4356.0	03 31 16.6	43 56 01.6	1.7	64.4	9.6 ± 0.9	10.2	4.5	3	224.5	11.7 ± 6.0	11.0	22.0	9.6	GK Per	CV, IP
SAXWFC J0336.8+0034.9 ^b	03 36 46.5	00 34 55.5	1.6	78.2	0.0 ± 0.0	0.0	0.0	3	251.7	12.9 ± 19.7	5.4	49.0	10.0	V711 Tau	RSCVn
SAXWFC J0355.4+3103.3	03 55 22.9	31 03 20.8	1.3	164.8	30.2 ± 0.9	38.8	13.1	25	1061.3	21.4 ± 5.0	10.3	35.0	16.0	4U 0352+309	Be HMXRB, XP
SAXWFC J0413.4+1026.6	04 13 23.8	10 26 34.0	2.0	50.3	7.9 ± 1.1	6.7	0.0	1	—	—	—	—	—	ABELL 478	CIG
SAXWFC J0433.2+0521.1	04 33 09.3	05 21 06.1	2.1	50.1	8.2 ± 1.1	7.3	1.0	2	89.5	6.3 ± 2.2	5.0	8.2	5.1	3C 120	AGN, Seyfert 1
SAXWFC J0452.0+4932.1	04 52 02.9	49 32 08.5	1.8	135.5	4.6 ± 0.5	8.1	0.0	5	365.5	4.4 ± 1.1	3.2	6.3	3.2	1RXS J045205.0+493248	AGN, Seyfert 1
SAXWFC J0458.2-7516.4	04 58 14.0	-75 16 21.3	1.8	70.5	8.6 ± 1.0	7.4	1.6	3	116.8	9.3 ± 1.0	0.6	11.0	8.6	1RXS J045816.3-751608	star KIII
SAXWFC J0507.7+6738.3	05 07 44.5	67 38 16.4	1.7	208.0	3.9 ± 0.4	9.5	2.5	6	977.3	4.3 ± 1.6	2.6	7.9	3.8	1ES0502+675	AGN, BL Lac
SAXWFC J0514.1-4002.3	05 14 07.3	-40 02 19.3	1.3	46.6	34.9 ± 0.9	56.4	16.5	38	1567.5	11.0 ± 7.6	55.2	35.0	4.5	4U 0513-40	LMXRB, GIC, B
SAXWFC J0520.4-7157.6	05 20 24.4	-71 57 37.0	1.3	191.8	53.1 ± 1.1	118.5	13.0	269	16241.2	51.3 ± 10.0	15.1	102.0	27.0	LMC X-2	LMXRB
SAXWFC J0528.8-6527.0	05 28 46.6	-65 27 00.3	1.5	9.1	45.9 ± 3.2	13.4	2.6	6	303.2	9.4 ± 15.1	31.0	46.0	5.6	AB Dor	star KIIIV
SAXWFC J0529.2-3248.3	05 29 13.6	-32 48 18.3	1.8	82.3	7.3 ± 1.0	6.8	2.5	4	190.6	6.8 ± 1.0	1.3	7.7	5.5	TV Col	CV
SAXWFC J0532.9-6622.0	05 32 54.5	-66 22 00.1	1.3	61.5	16.8 ± 0.6	31.4	30.0	67	3706.5	15.4 ± 7.9	31.9	49.0	3.6	LMC X-4	SG HMXRB, XP
SAXWFC J0534.5+2201.2	05 34 30.9	22 01 09.4	1.3	23.5	2071.4 ± 41.5	751.4	155.0	152	4635.1	2113.8 ± 89.4	5.1	2571.0	1680.7	Crab Nebula	SNR filled-center
SAXWFC J0535.3-0525.4	05 35 18.7	-05 25 21.0	1.9	30.5	6.4 ± 1.1	5.9	0.0	1	—	—	—	—	—	AXS J053514-0523	unclassified
SAXWFC J0539.0-6405.2	05 38 57.5	-64 05 15.0	1.3	78.3	66.8 ± 1.4	122.4	5.1	167	7781.3	36.8 ± 25.2	192.5	215.0	5.9	LMC X-3	HMXRB, BH
SAXWFC J0539.6-6944.7	05 39 38.0	-69 44 44.1	1.3	69.5	49.6 ± 1.1	99.7	5.8	209	11334.9	39.7 ± 7.8	14.2	77.0	24.0	LMC X-1	SG HMXRB, BH
SAXWFC J0540.1-6919.2	05 40 05.3	-69 19 14.8	1.7	30.7	5.9 ± 0.8	7.8	2.9	6	720.7	4.1 ± 1.0	2.1	5.9	3.3	PSR J0540-6919	XP
SAXWFC J0542.8+6051.2	05 42 50.5	60 51 11.1	1.8	161.6	4.6 ± 0.6	7.9	3.4	4	611.7	4.3 ± 0.4	0.5	4.6	3.8	BY Cam	CV, AM Her
SAXWFC J0550.6-3216.2	05 50 34.9	-32 16 11.2	1.7	91.7	4.5 ± 0.5	9.0	1.6	2	172.2	4.7 ± 0.8	1.2	5.6	4.5	PKS 0548-322	AGN, BL Lac
SAXWFC J0552.1-0727.9 ^f	05 52 06.3	-07 27 51.1	2.0	44.8	0.0 ± 0.0	0.0	0.0	3	302.4	13.2 ± 24.6	6.2	57.0	3.0	NGC 2110	AGN, Seyfert 2
SAXWFC J0552.5+5928.9	05 52 28.2	59 28 53.0	2.0	135.5	4.2 ± 0.6	6.7	1.1	1	—	—	—	—	—	IRAS 05480+5927	AGN, Seyfert 1
SAXWFC J0554.9+4625.6	05 54 55.2	46 25 33.6	1.5	125.6	6.9 ± 0.5	14.7	5.2	7	377.2	6.9 ± 1.8	3.6	11.0	5.7	MCG+08-11-011	AGN, Seyfert 1
SAXWFC J0617.1+0908.9	06 17 07.4	09 08 51.3	1.3	80.2	129.9 ± 2.9	105.3	27.6	95	1908.9	83.1 ± 46.7	53.6	360.0	37.0	4U 0614+091	LMXRB, B
SAXWFC J0620.8+1327.0	06 20 49.9	13 27 01.0	2.3	97.8	43.0 ± 6.9	5.5	1.8	1	326.9	—	—	—	—	WD 0618+134	WD, DA
SAXWFC J0637.0-0522.3	06 36 57.8	-05 22 16.3	1.7	42.3	9.7 ± 0.7	13.0	1.8	1	—	—	—	—	—	1RXS J063656.7-052104	unclassified
SAXWFC J0714.3-3625.4	07 14 16.4	-36 25 23.1	2.1	23.8	9.7 ± 1.3	7.1	1.6	1	—	—	—	—	—	1RXS J071413.7-362533	unclassified
SAXWFC J0728.9-2606.4	07 28 51.6	-26 06 24.8	1.5	62.8	17.0 ± 1.5	10.4	5.2	5	248.2	10.2 ± 4.4	10.4	17.0	7.3	4U 0728-25	Be HMXRB, XP
SAXWFC J0743.2+2853.1	07 43 12.8	28 53 06.0	2.0	99.5	5.6 ± 0.8	6.8	0.0	1	—	—	—	—	—	Sig Gem	RSCVn
SAXWFC J0747.5-1917.6	07 47 30.4	-19 17 35.1	1.7	46.7	6.3 ± 0.7	9.5	0.0	4	294.5	6.2 ± 0.4	0.2	6.7	5.8	PKS 0745-19	CIG

Table 3. BeppoSAX WFC X-ray Source Catalogue

WFC Name	RA(2000) hh mm ss	DEC(2000) o ' "	$\sigma_{ERR,RAD}$ arcmin	EXPO ks	$^bFlux_{2-10}$	$^cSNR1$	$^dSNR2$	N_{det}	EXPO _{total} ks	$^e<FLUX>_{2-10}$	χ^2_{flux}	$^fFlux_{MAX}$	$^fFlux_{MIN}$	Name	Class
SAXWFC J0748.5-6745.2	07 48 28.8	-67 45 14.7	1.3	56.6	43.2 ± 1.2	49.1	11.8	180	9893.5	15.8 ± 7.4	14.6	63.0	8.3	EXO 0748-676	LMXRB, B, E
SAXWFC J0749.1-0551.3	07 49 06.7	-05 51 16.2	2.2	42.5	51.7 ± 7.2	6.0	6.5	1	—	—	—	—	—	RX J0749.1-0549	CV, AM Her
SAXWFC J0812.5-3114.3	08 12 29.4	-31 14 17.8	1.5	47.3	11.7 ± 0.6	19.1	5.8	10	330.2	12.1 ± 10.8	31.5	42.0	5.2	RX J0812.4-3114	Be HMXRB, XP,
SAXWFC J0823.2-4250.1 ^h	08 23 11.7	-42 50 04.2	1.5	41.7	0.0 ± 0.0	0.0	0.0	28	1274.8	5.7 ± 5.4	2.1	19.0	4.8	1ES 0821-426	unclassified
SAXWFC J0835.3-4510.8	08 35 19.3	-45 10 50.5	1.6	42.5	6.7 ± 0.7	10.0	5.1	18	1086.5	7.3 ± 1.2	1.2	11.0	5.5	Vela Pulsar	SNR, XP, RP
SAXWFC J0841.4+7054.8	08 41 21.7	70 54 48.2	1.9	60.5	6.4 ± 0.8	8.1	2.0	10	924.9	4.9 ± 1.0	1.6	6.6	4.1	4C 71.07	QSO
SAXWFC J0902.1-4032.8	09 02 04.2	-40 32 46.6	1.3	102.0	187.0 ± 3.7	221.0	86.5	129	4770.0	82.5 ± 76.4	483.7	350.0	14.0	Vela X-1	SG HMXRB, XP,
SAXWFC J0920.4-5512.4	09 20 26.1	-55 12 24.1	1.3	6.0	174.3 ± 3.8	107.0	38.0	109	5273.6	14.4 ± 16.9	55.7	170.0	4.0	2S 0918-549	LMXRB, B
SAXWFC J0922.6-6317.6	09 22 38.4	-63 17 35.1	1.5	44.9	8.4 ± 0.7	11.0	2.6	21	1473.7	6.5 ± 2.4	8.2	12.0	3.6	2S 0921-63	LMXRB, E
SAXWFC J0923.7+2253.8	09 23 39.7	22 53 46.3	2.3	26.1	9.8 ± 1.2	7.4	0.0	1	—	—	—	—	—	RX J0923.7+2254	AGN, Seyfert 1
SAXWFC J0924.3-3142.4	09 24 17.7	-31 42 24.8	1.6	74.7	9.7 ± 0.6	14.5	0.0	32	1456.1	12.3 ± 4.3	6.7	26.0	8.5	1RXS J092418.0-314212	unclassified
SAXWFC J0934.1+6251.9	09 34 06.4	62 51 55.8	2.1	17.2	13.2 ± 1.8	6.8	2.8	1	—	—	—	—	—	FF UMa	star G5
SAXWFC J0945.6-1419.7 ⁱ	09 45 35.0	-14 19 44.0	1.9	66.9	7.0 ± 1.1	6.0	4.5	1	156.7	—	—	—	—	NGC 2992	AGN, Seyfert 1
SAXWFC J0947.6-3056.7	09 47 35.3	-30 56 43.8	1.9	76.8	10.4 ± 0.6	16.2	5.3	10	623.3	9.6 ± 2.6	8.9	15.0	6.8	MCG-5-23-16	AGN, Seyfert 2
SAXWFC J0955.5+6903.8	09 55 31.8	69 03 50.0	1.9	87.2	3.1 ± 0.4	7.2	3.6	4	356.1	3.8 ± 1.5	4.0	6.4	3.1	NGC 3031	AGN, Seyfert 1
SAXWFC J0955.6+6940.6	09 55 38.6	69 40 33.2	2.0	48.5	4.7 ± 0.4	10.4	2.8	3	217.6	4.8 ± 0.7	1.7	5.8	4.4	NGC 3034	Galaxy
SAXWFC J1009.7-5817.6	10 09 43.8	-58 17 35.5	1.3	87.2	33.0 ± 1.0	41.7	13.3	4	335.0	13.1 ± 11.7	172.4	33.0	7.8	GRO J1008-57	Be HMXRB, XP,
SAXWFC J1037.6-5647.6	10 37 34.8	-56 47 33.3	1.8	83.5	5.9 ± 0.5	11.6	6.8	3	368.8	6.4 ± 0.9	1.5	7.7	5.9	RX J1037.5-5647	Be HMXRB, XP
SAXWFC J1045.1-5941.1	10 45 03.2	-59 41 08.5	1.4	54.5	15.7 ± 0.7	23.1	6.9	42	2879.5	9.3 ± 4.1	11.1	21.0	5.4	Eta Car	LBV
SAXWFC J1052.6-6216.0	10 52 33.3	-62 15 59.4	1.7	39.3	17.6 ± 1.6	10.6	9.9	1	—	—	—	—	—	2E 1050.8-6200	unclassified
SAXWFC J1104.4+3812.0	11 04 23.0	38 12 00.3	1.3	71.4	53.0 ± 1.8	33.9	3.3	31	1517.1	21.2 ± 13.1	88.6	58.0	6.5	MKN 421	AGN, BL Lac
SAXWFC J1106.8+7233.6	11 06 50.5	72 33 34.2	1.4	84.7	9.1 ± 0.6	14.0	3.6	19	1244.8	5.9 ± 1.9	7.6	11.0	3.3	NGC 3516	AGN, Seyfert 1
SAXWFC J1118.2+4802.9	11 18 14.9	48 02 55.6	1.3	79.2	69.2 ± 1.5	80.7	19.0	9	357.4	66.1 ± 7.7	14.5	80.0	54.0	XTE J1118+480	LMXRB, BH
SAXWFC J1121.3-6037.5	11 21 15.9	-60 37 30.0	1.3	67.0	559.9 ± 11.0	545.7	209.8	152	8498.3	40.4 ± 170.5	736.5	610.0	8.0	Cen X-3	HMXRB, XP
SAXWFC J1124.5-5915.7	11 24 31.6	-59 15 44.6	1.8	147.8	4.2 ± 0.6	6.8	0.9	4	529.9	3.3 ± 4.4	7.8	12.0	2.5	SNR 292.0+01.8	SNR
SAXWFC J1138.9-3744.7	11 38 56.0	-37 44 41.2	1.7	167.4	8.4 ± 0.8	10.7	3.5	12	975.7	7.1 ± 2.0	4.6	11.0	4.2	NGC 3783	AGN, Seyfert 1
SAXWFC J1139.5-6523.4	11 39 28.1	-65 23 21.8	1.3	96.7	38.8 ± 1.0	60.1	11.9	10	876.2	14.0 ± 19.2	123.3	64.0	6.1	HD 101379	star G5III
SAXWFC J1141.5-6411.3	11 41 29.6	-64 11 18.2	2.3	130.0	6.0 ± 0.8	7.0	3.2	2	225.5	5.1 ± 1.2	2.6	6.0	4.3	V1033 Cen	CV, AM Her
SAXWFC J1143.6+7141.1	11 43 35.2	71 41 04.5	1.7	60.5	4.5 ± 0.6	7.8	0.0	3	201.1	4.6 ± 0.1	0.1	4.7	4.4	YY Dra	CV
SAXWFC J1143.8-6108.3	11 43 48.2	-61 08 17.8	1.7	45.3	14.7 ± 1.0	14.2	9.2	1	—	—	—	—	—	IGR J11435-6109	Be HMXRB, T
SAXWFC J1147.4-6156.5	11 47 21.5	-61 56 32.2	1.3	80.5	41.0 ± 1.3	35.7	23.5	71	5291.0	16.8 ± 6.8	19.0	54.0	6.8	1E 1145.1-6141	SG HMXRB, XP
SAXWFC J1148.0-6211.9	11 48 02.4	-62 11 51.3	1.3	17.9	108.8 ± 3.0	47.9	27.5	6	308.5	25.7 ± 38.4	365.6	110.0	11.0	4U 1145-619	Be HMXRB, XP
SAXWFC J1210.4+3924.4	12 10 25.2	39 24 26.2	1.3	66.5	12.7 ± 0.6	24.8	15.4	40	1666.5	14.8 ± 6.5	22.1	33.0	8.0	NGC 4151	AGN, Seyfert 1
SAXWFC J1213.2-6452.7	12 13 10.7	-64 52 44.7	1.3	96.6	25.7 ± 0.9	36.3	13.6	20	1099.7	18.5 ± 7.6	34.3	34.0	8.3	1ES 1210-64.6	unclassified
SAXWFC J1226.6-6245.7	12 26 36.1	-62 45 43.2	1.3	22.8	173.4 ± 3.6	175.5	76.6	105	6548.9	37.3 ± 53.7	314.2	340.0	8.1	GX 301-2	SG HMXRB, XP
SAXWFC J1229.1+0202.3	12 29 05.4	02 02 20.7	1.9	22.5	7.9 ± 0.9	8.1	0.6	3	32.3	9.8 ± 6.7	10.7	21.0	7.9	3C 273	AGN, BL Lac
SAXWFC J1248.9-4119.6	12 48 54.7	-41 19 35.4	1.6	97.6	5.8 ± 0.7	8.2	1.2	1	—	—	—	—	—	ABELL 3526	CIG
SAXWFC J1249.7-5905.1	12 49 42.0	-59 05 04.2	1.3	52.2	47.8 ± 1.2	61.7	13.4	50	3436.9	22.1 ± 15.5	72.2	73.0	9.1	4U 1246-58	LMXRB, B
SAXWFC J1252.4-2914.0	12 52 21.6	-29 13 58.8	1.5	32.7	10.7 ± 0.7	15.5	4.4	8	545.9	10.4 ± 1.4	1.1	14.0	9.4	EX Hya	CV
SAXWFC J1257.6-6917.0	12 57 35.3	-69 17 00.9	1.3	135.7	78.1 ± 1.6	116.7	12.5	178	12118.0	69.5 ± 12.4	18.3	130.0	38.0	4U 1254-690	LMXRB, B, D
SAXWFC J1259.5+2756.6	12 59 31.7	27 56 34.4	1.5	75.4	9.8 ± 1.1	8.5	1.2	1	—	—	—	—	—	Coma Cluster	CIG
SAXWFC J1325.5-4302.1	13 25 29.4	-43 02 03.1	1.4	148.8	20.5 ± 0.7	34.9	18.5	32	1863.7	23.2 ± 5.3	16.1	33.0	13.0	Cen A	AGN, Seyfert 2
SAXWFC J1326.7-6208.8	13 26 42.7	-62 08 47.0	1.4	124.3	36.8 ± 1.6	22.0	1.4	33	2724.2	14.7 ± 12.9	31.9	59.0	8.0	4U 1323-619	LMXRB, B, D
SAXWFC J1336.0-3417.1	13 35 57.2	-34 17 07.0	1.6	141.9	6.2 ± 0.5	11.0	3.9	1	—	—	—	—	—	MCG-06-30-015	AGN, Seyfert 1
SAXWFC J1347.4-3252.0	13 47 25.4	-32 51 59.0	1.6	140.8	7.0 ± 0.7	9.4	2.6	4	356.9	7.8 ± 0.8	0.9	9.0	7.0	ABELL 3571	CIG
SAXWFC J1348.8+2635.9	13 48 50.9	26 35 53.5	1.6	78.2	6.3 ± 0.7	8.2	3.1	4	148.9	6.8 ± 0.8	0.5	8.1	6.3	PKS 1346+26	AGN, RG
SAXWFC J1349.3-3019.1	13 49 17.6	-30 19 07.3	1.4	96.4	18.4 ± 0.9	18.9	5.9	14	780.2	14.5 ± 2.5	4.0	19.0	11.0	ESO 445-50	AGN, Seyfert 1
SAXWFC J1358.0-6443.7	13 58 03.0	-64 43 39.0	1.3	24.8	84.3 ± 2.5	41.2	15.8	7	269.6	55.9 ± 23.0	84.4	84.0	25.0	Cen X-2	LMXRB, BH, T
SAXWFC J1413.3-0311.7	14 13 15.4	-03 11 44.5	1.5	58.5	10.4 ± 0.6	19.0	6.0	8	429.8	9.2 ± 2.2	3.8	13.0	6.3	MKN 1376	AGN, Seyfert 2
SAXWFC J1421.2-6241.7	14 21 12.3	-62 41 42.0	1.5	33.2	17.0 ± 1.2	15.2	15.8	13	808.8	18.7 ± 9.2	17.0	37.0	9.5	4U 1416-62	Be HMXRB, XP,
SAXWFC J1428.6-5421.9	14 28 37.2	-54 21 56.8	1.5	107.5	32.3 ± 1.9	16.1	6.7	2	210.5	27.2 ± 6.5	13.4	32.0	23.0	SAX J1428.6-5422	T, unclassified
SAXWFC J1513.9-5908.8	15 13 55.3	-59 08 49.5	1.7	56.3	9.5 ± 1.5	6.2	6.0	1	—	—	—	—	—	PSR B1509-58	RP
SAXWFC J1520.6-5709.8	15 20 38.3	-57 09 40.4	1.3	58.0	2020.2 ± 40.1	537.5	190.9	95	5318.1	1563.6 ± 1072.0	631.4	5700.0	530.0	Cir X-1	XB, B, T
SAXWFC J1542.4-5222.7	15 42 26.1	-52 22 43.6	1.4	58.7	58.2 ± 3.0	20.1	28.7	20	1508.8	35.6 ± 14.6	13.8	72.0	22.0	4U 1538-522	SG HMXRB, XP

Verrecchia et al.: The BeppoSAX WFC source catalogue

Table 3. BeppoSAX WFC X-ray Source Catalogue

WFC Name	RA(2000) hh mm ss	DEC(2000) ° ' "	$\sigma_{ERR,RAD}$ arcmin	EXPO ks	$^b Flux_{2-10}$	$^c SNR1$	$^d SNR2$	N_{det}	EXPO _{total} ks	$^e <FLUX>_{2-10}$	χ^2_{flux}	$^f Flux_{MAX}$	$^f Flux_{MIN}$	Name	Class
SAXWFC J1544.0-5646.5	15 44 01.8	-56 46 32.5	1.4	152.5	16.1 ± 2.0	7.8	12.0	1	—	—	—	—	—	XTE J1543-568	Be HMXRB,
SAXWFC J1547.9-6234.6	15 47 55.5	-62 34 34.6	1.3	121.3	90.0 ± 2.3	64.6	32.1	86	5519.3	70.2 ± 20.4	19.1	150.0	33.0	4U 1543-624	LMXRB
SAXWFC J1550.9-5628.5	15 50 52.8	-56 28 27.8	1.3	35.4	5078.8 ± 99.3	305.0	40.8	20	1394.6	83.6 ± 2523.5	1770.5	6100.0	41.0	XTE J1550-564	LMXRB, BH
SAXWFC J1558.3+2712.7	15 58 19.4	27 12 43.9	1.6	80.0	6.5 ± 0.7	9.1	3.4	8	394.6	7.0 ± 2.5	6.8	11.0	4.1	ABELL 2142	CIG
SAXWFC J1601.1-6044.3	16 01 06.1	-60 44 20.4	1.4	60.4	40.7 ± 1.5	32.0	6.2	47	3400.5	41.3 ± 6.8	2.6	61.0	31.0	4U 1556-60	LMXRB
SAXWFC J1612.7-5224.6	16 12 39.5	-52 24 36.0	1.3	40.8	1146.4 ± 23.2	222.5	115.4	18	704.3	104.3 ± 322.3	280.8	1100.0	44.0	4U 1608-522	LMXRB, B,
SAXWFC J1619.9-1537.4	16 19 54.1	-15 37 25.3	1.3	6.9	21072.2 ± 414.7	570.6	111.1	99	1974.0	24754.9 ± 4116.0	48.6	40000.0	19000.0	Sco X-1	LMXRB
SAXWFC J1628.0-4912.0	16 28 00.6	-49 12 02.8	1.3	131.1	125.3 ± 3.4	50.4	27.3	54	2920.1	115.9 ± 17.1	5.0	170.0	77.0	4U 1624-490	LMXRB, D
SAXWFC J1628.6+3931.5	16 28 37.8	39 31 27.8	1.6	78.9	7.0 ± 0.8	8.6	2.8	16	761.1	6.0 ± 1.7	2.7	10.0	4.6	ABELL 2199	CIG
SAXWFC J1632.1-6727.7	16 32 07.8	-67 27 40.3	1.5	45.3	16.6 ± 0.8	20.0	11.8	66	4581.5	16.5 ± 3.2	3.4	27.0	9.8	4U 1627-673	LMXRB, XP
SAXWFC J1634.0-4722.8	16 33 59.0	-47 22 50.8	1.3	39.4	564.4 ± 12.1	98.7	55.6	9	646.2	124.4 ± 210.8	415.0	590.0	69.0	4U 1630-472	LMXRB, BH
SAXWFC J1638.1-6422.2	16 38 04.9	-64 22 13.8	1.7	68.9	9.8 ± 1.2	7.2	0.0	2	73.8	11.2 ± 7.3	9.8	20.0	9.8	IGR J16377-6423	CIG
SAXWFC J1641.0-5345.2	16 40 57.7	-53 45 11.5	1.3	51.7	634.3 ± 12.8	214.2	65.2	69	3285.2	32.9 ± 107.5	1332.5	700.0	6.4	4U 1636-536	LMXRB, B
SAXWFC J1645.8-4536.3	16 45 45.5	-45 36 18.7	1.3	56.1	960.3 ± 19.1	280.0	97.1	129	6769.2	25.0 ± 151.9	1588.0	1300.0	10.0	GX 340+0	LMXRB
SAXWFC J1653.8+3944.2	16 53 47.4	39 44 15.0	1.5	23.5	21.9 ± 0.9	28.1	11.0	70	3245.3	10.3 ± 8.2	43.8	42.0	4.2	MKN 501	AGN, BL La
SAXWFC J1654.0-3950.9	16 53 57.9	-39 50 52.0	1.3	14.6	6810.3 ± 133.3	504.5	167.6	42	1448.0	59.9 ± 1876.6	2401.7	7400.0	43.0	GRO J1655-40	LMXRB, BH
SAXWFC J1657.9+3521.1	16 57 51.6	35 21 03.2	1.3	72.3	98.7 ± 2.0	177.0	68.7	42	2061.1	18.4 ± 70.5	524.7	240.0	4.1	Her X-1	LMXRB, XP
SAXWFC J1700.7-4139.4	17 00 43.0	-41 39 21.2	1.6	72.8	112.7 ± 8.5	12.0	11.2	13	695.4	61.3 ± 28.0	14.7	120.0	30.0	OA0 1657-415	SG HMXRB
SAXWFC J1702.1-2957.9	17 02 05.9	-29 57 56.1	1.8	74.6	76.8 ± 4.7	15.9	6.6	25	1452.3	64.9 ± 7.5	2.0	77.0	49.0	XB 1658-298	LMXRB, B,
SAXWFC J1702.9-4847.5	17 02 51.1	-48 47 30.4	1.3	84.6	481.5 ± 10.1	100.9	3.7	19	736.1	123.2 ± 205.1	362.9	600.0	41.0	GX 339-4	LMXRB, BH
SAXWFC J1703.9-3750.9	17 03 54.5	-37 50 56.7	1.3	86.9	286.8 ± 7.9	48.8	76.7	96	4763.3	137.8 ± 81.5	80.2	440.0	46.0	4U 1700-377	SG HMXRB
SAXWFC J1704.2+7838.0	17 04 11.3	78 38 02.0	1.5	206.7	4.2 ± 0.5	8.6	2.4	11	977.4	4.4 ± 6.8	6.2	21.0	3.0	NGC 6331	Galaxy
SAXWFC J1705.7-3625.1	17 05 44.4	-36 25 03.3	1.3	17.5	1393.4 ± 27.9	355.2	103.6	154	6154.0	36.2 ± 247.4	2058.0	2100.0	14.0	GX 349+2	LMXRB
SAXWFC J1706.2-4302.0	17 06 11.9	-43 01 58.0	1.3	37.2	202.8 ± 5.8	46.7	37.2	81	4513.3	89.7 ± 45.0	36.8	260.0	43.0	4U 1702-429	LMXRB, B
SAXWFC J1706.6+2357.5	17 06 35.9	23 57 27.0	1.7	5.1	26.8 ± 2.5	10.2	2.7	5	87.7	16.0 ± 6.4	12.8	27.0	11.0	4U 1700+24	LMXRB
SAXWFC J1708.9-4406.9	17 08 52.1	-44 06 53.2	1.3	18.3	698.9 ± 14.2	160.3	54.4	114	5401.1	8.5 ± 227.8	968.3	920.0	2.0	4U 1705-440	LMXRB, B
SAXWFC J1711.6-3807.3	17 11 34.2	-38 07 17.7	1.3	86.2	108.2 ± 4.4	26.7	27.6	5	320.5	99.2 ± 12.4	4.7	110.0	79.0	SAX J1711.6-3808	LMXRB, BH
SAXWFC J1712.4-4050.7	17 12 21.3	-40 50 43.0	1.4	86.6	94.3 ± 3.8	27.6	9.8	33	1744.4	71.8 ± 13.7	5.1	100.0	51.0	4U 1708-40	LMXRB, B
SAXWFC J1712.6-3739.5	17 12 35.4	-37 39 31.3	1.4	34.5	76.6 ± 5.2	13.8	2.2	8	354.2	56.5 ± 11.1	5.8	77.0	43.0	SAX J1712.6-3739	LMXRB, B,
SAXWFC J1714.3-3402.8	17 14 18.5	-34 02 48.8	1.4	40.5	87.7 ± 4.8	18.6	7.7	12	667.9	79.3 ± 14.8	5.1	100.0	62.0	4U 1711-34	LMXRB, B,
SAXWFC J1716.0-3851.9	17 15 59.8	-38 51 53.6	1.4	17.7	62.7 ± 4.2	15.4	3.9	43	1993.8	62.8 ± 13.6	3.6	100.0	40.0	XTE J1716-389	SG HMXRB
SAXWFC J1723.6-3739.4	17 23 37.2	-37 39 24.1	1.3	36.0	125.1 ± 4.4	32.9	17.0	9	511.8	123.3 ± 28.6	23.6	150.0	74.0	XTE J1723-376	LMXRB, B,
SAXWFC J1727.5-3048.1	17 27 29.4	-30 48 04.6	1.4	50.4	76.5 ± 4.7	16.5	12.3	74	3990.1	51.4 ± 20.2	6.4	170.0	34.0	4U 1724-307	LMXRB, GLO
SAXWFC J1728.6+5901.5	17 28 34.0	59 01 29.2	1.4	15.1	38.9 ± 1.5	27.7	5.1	5	166.0	15.0 ± 14.2	157.3	39.0	6.8	GR Dra	Star G0
SAXWFC J1731.7-1657.6	17 31 42.8	-16 57 38.8	1.3	35.6	473.3 ± 10.4	81.9	26.5	163	5076.0	501.8 ± 76.5	11.1	980.0	350.0	GX 9+9	LMXRB
SAXWFC J1731.9-3350.2	17 31 55.1	-33 50 10.3	1.3	60.4	356.4 ± 8.2	79.5	64.9	135	5331.8	10.7 ± 75.3	534.5	440.0	1.5	GX 354-0	LMXRB, B
SAXWFC J1732.0-2444.0	17 32 01.1	-24 44 02.0	1.4	41.1	89.5 ± 5.4	16.8	41.4	24	1180.4	51.4 ± 17.9	9.8	89.0	33.0	GX 1+4	LMXRB, XP
SAXWFC J1732.5+7413.2	17 32 31.0	74 13 14.1	2.4	33.5	4.4 ± 0.5	7.8	0.0	1	—	—	—	—	—	DR Dra	RSCVn
SAXWFC J1733.4-3323.8	17 33 22.7	-33 23 47.7	1.3	55.5	479.3 ± 10.5	98.6	78.5	24	924.0	192.6 ± 120.8	211.8	480.0	47.0	Rapid Burster	LMXRB, GLO
SAXWFC J1734.2-2605.2	17 34 10.5	-26 05 13.5	1.3	126.0	373.2 ± 8.1	105.3	39.2	115	4500.5	23.6 ± 149.1	625.9	630.0	5.3	KS 1731-260	LMXRB, B,
SAXWFC J1738.2-2659.5	17 38 13.2	-26 59 27.6	1.5	15.0	58.0 ± 5.8	10.1	2.2	9	465.7	24.6 ± 12.4	6.8	58.0	18.0	SLX 1735-269	LMXRB, B
SAXWFC J1738.9-4426.8	17 38 56.7	-44 26 48.4	1.3	129.2	410.0 ± 8.2	167.9	59.2	115	5856.2	16.5 ± 85.4	1206.4	520.0	3.5	4U 1735-444	LMXRB, B
SAXWFC J1740.1-3103.0 ⁱ	17 40 07.6	-31 03 00.0	1.6	127.5	28.5 ± 4.8	5.8	8.5	6	704.9	24.1 ± 24.8	5.3	81.0	11.0	GRS 1737-310	LMXRB, T
SAXWFC J1740.7-2818.5	17 40 40.8	-28 18 30.2	1.6	17.5	123.7 ± 9.3	12.4	10.1	1	—	—	—	—	—	SLX 1737-282	LMXRB, B
SAXWFC J1742.7-2744.9	17 42 39.6	-27 44 51.7	1.3	151.1	360.7 ± 8.6	74.6	2.7	26	636.6	231.5 ± 79.2	96.5	370.0	80.0	GRS 1739-278	LMXRB, BH
SAXWFC J1743.8-2945.1	17 43 48.5	-29 45 03.9	1.7	60.8	35.5 ± 3.2	11.4	20.9	50	2914.5	30.2 ± 6.9	2.0	63.0	20.0	1E 1740.7-2942	LMXRB, BH
SAXWFC J1744.5-2844.6	17 44 31.6	-28 44 35.5	1.3	126.8	353.2 ± 8.4	76.2	154.8	20	1181.2	100.7 ± 174.7	221.1	670.0	32.0	GRO J1744-28	LMXRB, XP
SAXWFC J1744.9-2921.2	17 44 52.0	-29 21 10.8	2.0	55.2	26.5 ± 4.2	6.4	8.6	3	174.7	29.7 ± 4.4	1.5	34.0	27.0	KS 1741-293	LMXRB, B
SAXWFC J1746.0-2931.0	17 46 01.6	-29 30 59.0	1.4	56.5	69.9 ± 3.1	24.6	20.4	57	3236.2	39.8 ± 16.1	16.2	84.0	23.0	1A 1742-294	LMXRB, B
SAXWFC J1747.0-2853.3	17 46 59.2	-28 53 17.8	1.3	79.2	242.2 ± 6.2	61.7	31.4	16	754.5	75.3 ± 69.7	141.7	240.0	27.0	SAX J1747.0-2853	LMXRB, B,
SAXWFC J1747.5-3001.5	17 47 28.1	-30 01 28.2	1.3	60.2	71.6 ± 4.0	19.2	14.0	58	3353.4	46.3 ± 12.3	8.1	77.0	26.0	SLX 1744-300	LMXRB, B
SAXWFC J1747.9-3009.0	17 47 54.1	-30 08 58.5	1.8	6.8	107.6 ± 14.9	6.6	1.0	0	—	—	—	—	—	1RXS J174735.7-300953	unclassified
SAXWFC J1747.9-2633.3	17 47 56.8	-26 33 16.5	1.3	63.4	867.3 ± 17.5	241.2	114.6	148	5428.6	31.6 ± 175.7	1634.6	1500.0	6.7	GX 3+1	LMXRB, B
SAXWFC J1748.1-2446.1	17 48 06.4	-24 46 08.7	1.3	41.2	866.2 ± 18.0	149.8	115.7	13	587.2	240.7 ± 262.2	644.9	890.0	37.0	EXO 1745-248	LMXRB, GLO
SAXWFC J1748.9-2021.3	17 48 53.8	-20 21 17.6	1.3	20.9	336.5 ± 8.5	62.1	25.1	5	156.3	133.2 ± 144.3	386.5	340.0	38.0	4U 1745-203	LMXRB, GLO

Table 3. BeppoSAX WFC X-ray Source Catalogue

WFC Name	RA(2000) hh mm ss	DEC(2000) o ' "	$\sigma_{ERR,RAD}$ arcmin	EXPO ks	$^b Flux_{2-10}$	$^c SNR1$	$^d SNR2$	N_{det}	EXPO _{total} ks	$^e <FLUX>_{2-10}$	χ^2_{flux}	$f Flux_{MAX}$	$f Flux_{MIN}$	Name	Class
SAXWFC J1750.2-3703.1	17 50 09.4	-37 03 07.5	1.4	40.7	99.8 ± 4.4	24.6	15.3	71	3785.7	82.6 ± 17.8	9.0	130.0	46.0	4U 1746-370	LMXRB, GIC, B
SAXWFC J1750.4-2902.6	17 50 23.5	-29 02 35.1	1.4	127.0	74.3 ± 4.0	19.6	8.9	3	198.8	81.3 ± 45.9	35.8	150.0	72.0	SAX J1750.8-2900	LMXRB, B, T
SAXWFC J1750.8-3116.5	17 50 47.5	-31 16 28.2	1.4	42.3	58.0 ± 3.5	17.1	12.5	18	977.4	47.6 ± 14.1	11.5	66.0	24.0	GRS 1747-312	LMXRB, GIC, B
SAXWFC J1751.2-3037.7	17 51 10.0	-30 37 41.1	1.4	15.9	89.6 ± 4.6	20.7	19.1	1	—	—	—	—	—	XTE J1751-305	XB, XP
SAXWFC J1755.5-3229.7	17 55 32.8	-32 29 42.7	1.5	22.9	66.8 ± 4.4	15.1	0.5	2	90.9	58.1 ± 10.2	6.6	67.0	52.0	XTE J1755-324	LMXRB, BH, T
SAXWFC J1801.1-2504.9	18 01 05.5	-25 04 56.6	1.3	64.6	2044.4 ± 40.4	411.3	105.4	147	5325.7	46.7 ± 378.3	2350.5	4700.0	18.0	GX 5-1	LMXRB
SAXWFC J1801.2-2544.4	18 01 13.1	-25 44 25.0	1.7	67.3	49.7 ± 4.1	12.1	11.2	20	1322.8	45.1 ± 9.2	2.2	66.0	34.0	GRS 1758-258	LMXRB, BH
SAXWFC J1801.5-2031.6	18 01 31.9	-20 31 34.3	1.3	31.0	1151.8 ± 23.6	220.5	122.7	143	5084.5	135.1 ± 199.0	1921.3	2500.0	8.9	GX 9+1	LMXRB
SAXWFC J1805.5-2031.1	18 05 30.5	-20 31 06.2	1.6	20.6	84.6 ± 6.7	12.1	1.1	3	158.6	66.3 ± 16.2	5.8	85.0	56.0	SAX J1805.5-2031	T
SAXWFC J1806.9-2434.7	18 06 51.3	-24 34 44.0	2.3	58.1	31.7 ± 5.0	6.1	6.8	1	—	—	—	—	—	2S 1803-245	LMXRB, B, T
SAXWFC J1808.4-3659.0	18 08 26.1	-36 59 01.3	1.3	7.9	177.9 ± 5.2	43.7	14.5	14	118.3	159.7 ± 56.4	31.8	230.0	57.0	SAX J1808.4-3658	LMXRB, XP, B,
SAXWFC J1810.8-2609.0	18 10 48.8	-26 09 02.5	2.0	57.7	40.0 ± 6.3	6.1	4.1	1	—	—	—	—	—	SAX J1810.8-2609	LMXRB, B, T
SAXWFC J1814.5-1708.8	18 14 28.1	-17 08 48.8	1.3	58.3	831.6 ± 17.3	146.9	41.6	161	4931.2	737.2 ± 96.9	162.2	16.2	530.0	GX 13+1	LMXRB, B
SAXWFC J1815.1-1205.6	18 15 08.2	-12 05 38.0	1.7	35.5	40.2 ± 4.2	9.1	3.8	9	252.1	37.4 ± 5.6	1.3	48.0	30.0	4U 1812-12	LMXRB, B
SAXWFC J1816.0-1402.6	18 16 01.0	-14 02 34.8	1.3	32.6	3580.1 ± 70.7	300.0	85.3	175	5293.6	1371.1 ± 217.3	24.0	3600.0	1100.0	GX 17+2	LMXRB, B
SAXWFC J1816.4+4953.1	18 16 21.0	49 53 04.2	1.4	224.8	14.3 ± 0.6	22.5	5.6	71	4007.7	10.3 ± 3.5	14.0	20.0	3.6	AM Her	CV, AM Her
SAXWFC J1819.4-2524.0	18 19 21.9	-25 24 00.3	1.5	69.3	72.9 ± 8.9	7.8	8.1	1	—	—	—	—	—	SAX J1819.3-2525	LMXRB, BH, T
SAXWFC J1821.9+6419.6	18 21 53.4	64 19 36.4	1.7	35.7	4.5 ± 0.6	7.0	1.7	3	202.9	3.1 ± 1.1	5.4	4.5	2.4	QSO B1821+643	AGN, Seyfert 1
SAXWFC J1823.7-3021.6	18 23 42.1	-30 21 38.8	1.3	312.3	818.0 ± 16.5	168.2	75.3	134	5360.6	27.6 ± 207.6	1338.6	940.0	4.3	4U 1820-303	LMXRB, GIC, B
SAXWFC J1825.2-0001.0	18 25 14.2	-00 00 58.6	1.4	15.5	74.5 ± 2.3	41.1	9.9	69	1318.6	67.1 ± 12.8	6.8	120.0	45.0	4U 1822-000	LMXRB
SAXWFC J1825.8-3707.0	18 25 46.0	-37 06 58.3	1.5	12.9	51.3 ± 2.5	20.2	11.2	61	3140.8	50.0 ± 7.8	1.9	72.0	39.0	4U 1822-371	LMXRB, E
SAXWFC J1829.6-2347.7	18 29 34.5	-23 47 39.4	1.4	36.5	82.7 ± 3.8	21.0	8.7	58	2554.4	73.7 ± 15.6	3.3	120.0	54.0	GS 1826-238	LMXRB, B
SAXWFC J1833.9+5143.5	18 33 53.2	51 43 28.2	1.5	25.8	11.4 ± 0.9	13.1	2.3	1	—	—	—	—	—	BY Dra	BY Dra
SAXWFC J1835.0+3241.6	18 35 01.8	32 41 37.6	1.5	128.1	5.5 ± 0.5	11.4	2.0	4	392.8	6.0 ± 0.5	0.8	6.7	5.5	3C 382	AGN, Seyfert 1
SAXWFC J1835.6-3258.2	18 35 38.1	-32 58 14.5	1.7	35.2	27.2 ± 3.5	7.5	4.7	3	76.0	21.9 ± 4.0	1.7	27.0	19.0	XB 1832-330	LMXRB, GIC, B
SAXWFC J1839.9+0501.9	18 39 54.7	05 01 51.2	1.3	46.5	530.8 ± 10.7	372.5	88.4	94	1830.7	480.0 ± 100.3	36.8	1100.0	240.0	Ser X-1	LMXRB, B
SAXWFC J1842.2+7945.6	18 42 13.2	79 45 38.1	1.6	62.1	5.6 ± 0.6	9.7	2.6	17	1691.6	4.0 ± 1.4	4.1	8.5	2.5	3C 390.3	AGN, Seyfert 1
SAXWFC J1845.6+0051.3	18 45 34.4	00 51 19.8	1.4	24.0	56.2 ± 2.2	28.7	32.4	11	209.4	43.3 ± 14.2	42.3	64.0	13.0	GS 1843+009	Be HMXRB, XP,
SAXWFC J1848.2-0225.9	18 48 14.4	-02 25 54.8	1.4	15.2	24.7 ± 2.4	10.3	17.7	4	114.2	27.2 ± 5.0	1.7	36.0	25.0	GS 1843-02	Be HMXRB, XP,
SAXWFC J1848.3-0310.6	18 48 18.0	-03 10 38.2	1.5	46.6	22.6 ± 2.4	9.3	7.4	0	—	—	—	—	—	IGR J18483-0311	unclassified
SAXWFC J1853.1-0844.0	18 53 03.5	-08 44 00.6	2.4	8.2	16.1 ± 2.1	7.3	1.7	1	—	—	—	—	—	4U 1850-087	LMXRB, GIC, B
SAXWFC J1855.5-0237.1 ⁱ	18 55 30.8	-02 37 08.7	2.2	43.6	7.4 ± 2.5	2.8	5.9	8	335.7	6.5 ± 8.6	2.7	29.0	2.3	XTE J1855-026	SG HMXRB, XP
SAXWFC J1856.8+0518.4	18 56 45.0	05 18 23.4	2.7	6.3	183.1 ± 5.8	38.9	2.3	11	160.3	120.7 ± 43.8	45.7	180.0	50.0	XTE J1856+053	LMXRB, BH, T
SAXWFC J1858.7+2240.0	18 58 42.9	22 39 57.2	1.3	81.0	1416.1 ± 28.9	133.7	62.0	5	357.9	13.6 ± 781.7	1140.8	1500.0	8.2	XTE J1859+226	LMXRB, BH, T
SAXWFC J1908.9+0922.3	19 08 51.9	09 22 15.9	1.3	8.4	86.4 ± 3.4	27.3	0.7	3	18.5	76.2 ± 10.8	7.9	86.0	66.0	XTE J1908+094	LMXRB, BH, T
SAXWFC J1909.6+0949.8	19 09 36.2	09 49 49.4	1.4	29.1	46.1 ± 2.7	16.7	10.9	19	605.7	22.4 ± 8.4	11.2	46.0	14.0	4U 1907+097	SG HMXRB, XP
SAXWFC J1910.8+0735.6	19 10 50.7	07 35 33.7	1.7	29.1	15.6 ± 1.7	9.4	7.7	8	319.3	16.7 ± 6.1	4.6	32.0	14.0	4U 1908+075	HMXRB, XP
SAXWFC J1911.3+0035.5	19 11 16.5	00 35 27.9	1.3	48.7	1602.9 ± 31.4	177.0	69.5	10	321.3	274.7 ± 497.7	662.9	1600.0	86.0	Aql X-1	LMXRB, B, T
SAXWFC J1911.8+0459.8	19 11 49.5	04 59 48.4	1.5	36.3	20.0 ± 2.1	9.1	2.8	5	168.5	19.4 ± 18.5	5.7	60.0	18.0	SS 433	SG HMXRB
SAXWFC J1915.2+1057.4	19 15 11.0	10 57 21.6	1.3	11.8	2565.9 ± 50.5	452.7	134.1	93	2582.0	889.1 ± 957.3	603.0	4300.0	460.0	GRS 1915+105	LMXRB, BH, T
SAXWFC J1918.8-0514.9	19 18 45.4	-05 14 51.3	1.4	36.3	36.8 ± 1.6	25.4	10.2	21	533.3	36.7 ± 17.1	33.9	77.0	16.0	XB 1916-053	LMXRB, B, D
SAXWFC J1921.1-5841.2	19 21 08.9	-58 41 09.6	1.6	72.1	5.8 ± 0.7	7.9	2.7	1	—	—	—	—	—	ESO 141-G55	AGN, Seyfert 1
SAXWFC J1921.2+4400.1	19 21 14.4	44 00 06.8	1.6	31.6	11.5 ± 1.3	8.2	0.8	9	493.0	8.9 ± 4.2	3.4	21.0	7.2	ABELL 2319	CIG
SAXWFC J1924.4+5014.5 ^j	19 24 26.9	50 14 28.6	1.8	68.7	0.0 ± 0.0	0.0	0.0	0	—	—	—	—	—	CH Cyg	Sy
SAXWFC J1928.5-3507.9	19 28 30.6	-35 07 51.2	1.5	42.3	17.6 ± 1.4	12.3	5.6	1	—	—	—	—	—	HD 182928	G5V star
SAXWFC J1945.7+2721.1	19 45 40.0	27 21 21.9	1.3	34.5	98.3 ± 2.4	69.1	55.4	21	1161.9	35.7 ± 30.7	168.4	100.0	11.0	XTE J1946+274	Be HMXRB, XP,
SAXWFC J1949.6+3011.8	19 49 34.2	30 11 47.0	1.3	48.6	165.9 ± 3.8	74.3	51.8	9	434.7	37.7 ± 56.6	204.1	170.0	24.0	KS 1947+300	Be HMXRB, XP,
SAXWFC J1955.7+3205.3	19 55 40.8	32 05 16.4	1.3	83.3	64.3 ± 1.7	58.8	37.7	35	1371.5	29.1 ± 21.4	80.8	110.0	12.0	4U 1954+319	HMXRB
SAXWFC J1958.4+3513.1	19 58 23.9	35 13 03.7	1.3	86.4	1162.9 ± 23.0	522.7	105.3	163	5471.8	586.3 ± 434.7	252.54	2300.0	170.0	Cyg X-1	SG HMXRB, BH
SAXWFC J1959.4+1141.8	19 59 22.2	11 41 46.3	1.3	52.5	46.5 ± 1.3	48.8	3.0	70	1850.1	64.7 ± 28.1	40.3	160.0	40.0	4U 1957+115	LMXRB, BH
SAXWFC J1959.4+4044.0	19 59 25.1	40 44 01.6	1.6	57.9	10.5 ± 1.4	7.2	2.9	2	109.9	9.7 ± 1.0	0.5	10.0	9.0	Cygnus A	QSO
SAXWFC J2000.0+6508.8	20 00 01.0	65 08 45.2	1.4	23.5	23.6 ± 0.9	28.7	5.7	42	3164.2	9.6 ± 4.4	18.8	25.0	4.6	1ES 1959+650	AGN, BL Lac
SAXWFC J2008.6-6107.4	20 08 35.7	-60 24 33.8	2.1	67.6	18.0 ± 2.5	6.3	0.0	1	—	—	—	—	—	1RXS J200905.6-602537	WD
SAXWFC J2008.6-6107.4	20 08 36.2	-61 07 23.5	2.0	91.5	4.9 ± 0.7	7.0	1.8	1	—	—	—	—	—	NGC 6860	AGN, Seyfert 1
SAXWFC J2009.4-4850.0	20 09 24.1	-48 50 00.2	1.3	70.4	32.9 ± 1.1	36.3	9.0	22	1045.1	16.2 ± 9.0	77.3	35.0	7.7	PKS 2005-489	AGN, BL Lac

Verrecchia et al.: The BeppoSAX WFC source catalogue

Table 3. BeppoSAX WFC X-ray Source Catalogue

WFC Name	RA(2000) hh mm ss	DEC(2000) ° ' "	$\sigma_{ERR,RAD}$ arcmin	EXPO ks	$^b Flux_{2-10}$	$^c SNR1$	$^d SNR2$	N_{det}	EXPO _{total} ks	$^e <FLUX>_{2-10}$	χ^2_{flux}	$^f Flux_{MAX}$	$^f Flux_{MIN}$	Name	Class
SAXWFC J2012.7+3811.0	20 12 39.3	38 10 57.7	1.3	3.3	289.2 ± 6.8	74.3	0.0	7	200.9	41.4 ± 141.0	573.8	310.0	16.0	XTE J2012+381	LMXRB, BH, T
SAXWFC J2012.7-5650.1	20 12 39.9	-56 50 05.2	1.6	74.5	4.6 ± 0.6	7.3	0.0	1	—	—	—	—	—	ABELL 3667	CIG
SAXWFC J2032.2+3738.2	20 32 13.6	37 38 14.6	1.4	27.9	31.8 ± 1.4	25.2	15.7	6	206.7	29.1 ± 6.6	8.8	37.0	21.0	EXO 2030+375	Be HMXRB, XP
SAXWFC J2032.5+4057.5	20 32 27.0	40 57 28.8	1.3	30.6	623.8 ± 12.4	311.2	67.3	170	6062.5	269.9 ± 206.9	421.3	890.0	130.0	Cyg X-3	HMXRB
SAXWFC J2044.2-1043.6	20 44 10.9	-10 43 35.0	1.8	85.8	6.1 ± 0.7	8.3	1.5	4	151.8	7.5 ± 4.9	6.2	17.0	6.1	MKN 509	AGN, Seyfert 1
SAXWFC J2058.8+4147.6	20 58 45.4	41 47 36.9	1.7	12.6	22.8 ± 2.6	8.8	7.4	3	59.9	18.7 ± 3.9	3.4	23.0	15.0	GRO J2058+42	Be HMXRB, XP
SAXWFC J2103.6+4545.9	21 03 38.1	45 45 55.8	1.4	104.4	46.0 ± 1.9	23.3	9.4	20	798.5	20.5 ± 11.6	29.5	58.0	7.2	SAX J2103.5+4545	Be HMXRB, XP
SAXWFC J2130.0+1209.7	21 29 59.5	12 09 44.6	1.3	20.1	51.9 ± 1.5	45.5	4.2	30	964.7	30.7 ± 11.1	39.8	54.0	17.0	4U 2127+119	LMXRB, B, GIC
SAXWFC J2132.0-3342.2	21 31 59.9	-33 42 10.4	2.2	39.5	4.5 ± 0.6	7.4	2.3	1	—	—	—	—	—	CTS 109	AGN, Seyfert 1
SAXWFC J2142.6+4335.8	21 42 37.3	43 35 50.2	1.7	53.6	24.6 ± 1.2	22.1	7.7	19	1125.6	20.0 ± 5.6	8.4	31.0	12.0	SS Cyg	CV, DN, U Gem
SAXWFC J2144.7+3818.8	21 44 42.0	38 18 46.0	1.3	22.8	1403.9 ± 28.0	576.0	106.4	145	4923.4	853.2 ± 335.9	307.9	1900.0	190.0	Cyg X-2	LMXRB, B, D
SAXWFC J2146.4+5842.7	21 46 21.6	-85 42 41.4	1.5	33.4	17.1 ± 1.8	9.6	2.1	0	—	—	—	—	—	1RXS J214641.1-854303	unclassified
SAXWFC J2148.8+8232.2	21 48 47.6	82 32 10.6	1.5	164.1	6.6 ± 0.6	11.7	6.7	0	—	—	—	—	—	SAXWFC J2148.8+8232	unclassified
SAXWFC J2158.8-3013.2	21 58 48.0	-30 13 09.8	1.9	46.3	7.6 ± 0.8	9.4	2.7	8	238.8	8.2 ± 6.2	14.1	24.0	5.6	PKS 2155-304	AGN, BL Lac
SAXWFC J2207.9+5431.9	22 07 56.9	54 31 52.3	1.4	82.5	26.6 ± 1.0	28.6	9.6	60	2873.9	13.9 ± 6.4	23.5	35.0	5.0	4U 2206+543	Be HMXRB, XP
SAXWFC J2208.8+4545.4	22 08 45.0	45 45 26.2	2.3	68.0	29.9 ± 4.2	6.1	2.5	1	—	—	—	—	—	AR Lac	CV, RSCVn
SAXWFC J2209.2-4713.3	22 09 14.3	-47 13 15.9	1.7	39.5	4.5 ± 0.7	6.1	0.0	1	—	—	—	—	—	NGC 7213	AGN, Seyfert 1
SAXWFC J2214.1+1242.6	22 14 06.5	12 42 34.5	1.8	66.6	5.4 ± 0.8	6.4	2.4	1	—	—	—	—	—	RU Peg	CV, DN, U Gem
SAXWFC J2239.2+6116.4	22 39 14.3	61 16 26.7	1.4	13.1	33.6 ± 1.6	21.6	18.1	1	—	—	—	—	—	SAX J2239.3+6116	Be HMXRB, XP
SAXWFC J2253.1+1649.3	22 53 05.4	16 49 15.6	1.5	43.4	24.2 ± 1.7	13.3	2.5	1	—	—	—	—	—	IM Peg	RSCVn
SAXWFC J2323.5+5848.1	23 23 27.6	58 48 03.9	1.3	138.2	109.8 ± 2.3	145.2	10.1	167	7876.7	103.4 ± 13.7	21.4	150.0	59.0	Cassiopeia A	SNR shell
SAXWFC J2331.9+1956.1	23 31 53.2	19 56 05.6	1.9	33.7	11.0 ± 1.4	7.0	0.0	1	—	—	—	—	—	EQ Peg	star M3V
SAXWFC J2359.2-3038.0	23 59 09.5	-30 37 59.8	2.2	45.9	5.3 ± 0.6	7.9	1.2	2	130.6	4.0 ± 1.6	7.4	5.3	3.1	1H 2354-315	AGN, BL Lac
SAXWFC J0234.3-4347.1 ^k	02 34 19.2	-43 47 07.4	1.9	113.7	3.9 ± 0.5	7.8	2.3	1	—	—	—	—	—	CC Eri	RSCVn
SAXWFC J0759.8-3845.0 ^k	07 59 45.0	-38 44 36.2	2.1	21.9	14.4 ± 2.1	6.0	0.0	1	—	—	—	—	—	IGR J07597-3842	AGN, Seyfert 1
SAXWFC J1510.9+0542.7 ^k	15 10 57.0	05 42 39.2	1.8	56.0	6.7 ± 0.9	7.3	0.0	2	100.8	7.1 ± 0.9	0.7	7.9	6.7	ABELL 2029	CIG
SAXWFC J1632.0-4751.5 ^k	16 32 02.0	-47 51 29.8	1.6	9.0	—	0.0	0.0	1	—	—	—	—	—	AX J1631.9-4752	HMXRB, XP
SAXWFC J1709.0-3623.2 ^k	17 09 00.6	-36 23 14.2	1.8	41.8	—	0.0	0.0	1	—	—	—	—	—	IGR J17091-3624	XB
SAXWFC J1709.5-2639.2 ^k	17 09 29.9	-26 39 11.8	2.1	93.3	52.6 ± 5.4	9.3	0.0	2	102.1	63.0 ± 5.4	27.3	129.0	52.6	1RXS J170930.2-263927	LMXRB
SAXWFC J1725.1+1153.8 ^k	17 25 07.2	11 53 45.2	2.1	13.7	15.6 ± 2.2	6.4	0.0	1	—	—	—	—	—	1H 1720+117	AGN, BL Lac
SAXWFC J1914.1+0951.7 ^k	19 14 07.7	09 51 42.8	2.0	6.7	—	0.0	0.0	1	—	—	—	—	—	IGR J19140+0951	HMXRB
SAXWFC J2055.4-0021.0 ^k	20 55 24.6	-00 20 59.2	1.9	50.0	4.96 ± 0.73	6.5	0.0	1	—	—	—	—	—	1RXS J205528.2-002123	unclassified
SAXWFC J2124.7+5058.3 ^k	21 24 40.8	50 58 17.7	2.0	81.0	6.96 ± 0.83	7.8	0.0	1	—	—	—	—	—	IGR J21247+5058	AGN, RG
SAXWFC J2355.2+2839.4 ^k	23 55 11.5	28 39 23.7	1.9	34.8	—	0.0	0.0	1	—	—	—	—	—	II Peg	RSCVn

^a the error radius is the mean 99% X, Y error added in quadrature to a systematic error of 1.3'

^b flux and error in units of $10^{-11} \text{ erg cm}^{-2} \text{ s}^{-1}$

^c signal-to-noise ratio in the 2-8 keV band

^d signal-to-noise ratio in the 8-19 keV band

^e the flux weighted mean among all source detections and its standard deviation in units of $10^{-11} \text{ erg cm}^{-2} \text{ s}^{-1}$

^f maximum and minimum fluxes in units of $10^{-11} \text{ erg cm}^{-2} \text{ s}^{-1}$

^g Sources classification: AGN, Active galactic nuclei; AM Her, AM Her type star; AXP, Anomalous X-ray pulsar; Be, B-emission star; B, Burster; BY Dra, BY Dra type star; CIG, Cluster of galaxies; CV, Cataclismic variable; D, dipper; DN, Dwarf nova; E, eclipsing; GIC, in globular cluster; HMXRB, High Mass X-ray Binaries; IP, Intermediate polar; LBV, LBV type star; LMXRB, Low Mass X-ray Binaries; P, Polar; QSO, Quasar; RG, radio galaxy; RP, radio pulsar; RSCVn, RS CVn type star; Seyfert 1 or 2, Seyfert galaxy type 1 or type 2; SNR, Supernova remnant; Sy, Symbiotic star; T, transient source; U Gem, U Gem type star; WD, White dwarf; XB, X-ray binary; XP, X-ray pulsar

^h detected with the highest snr in a softer band, 2–5.5 or 2–3.2 keV with respect to the catalogue reference 2–8 keV band

ⁱ detected with the highest snr in a harder band, 8–19, 5.5–11, 6–13 or 11–19 keV with respect to the catalogue reference 2–8 keV band

^j detected with the highest snr in a larger, 2–19 or 2–26 keV, band with respect to the catalogue reference 2–8 keV band

^k source detected in the first processing stage only